
World key to the species of Pyrenulaceae and Trypetheliaceae

André Aptroot¹

¹Laboratório de Botânica / Liquenologia, Instituto de Biociências Universidade Federal de Mato Grosso do Sul, Avenida Costa e Silva s/n Bairro Universitário, CEP 79070-900, Campo Grande, Mato Grosso do Sul, Brazil email: andreaptroot@gmail.com

The keys below treat all c. 750 species in the families Pyrenulaceae and Trypetheliaceae. They are a separate print of the keys published in this book:

FELIX SCHUMM & ANDRÉ APTROOT: Atlas of Pyrenulaceae and Trypetheliaceae, Volume 1, pp1-520, ISBN 978-3-7543-9685-8

They are published separately here online so as to give access to a wide audience, also to those who do not buy the book.

General key to the families and some genera of tropical pyrenocarpous lichens

(specimens without ascospores or with only conidia can rarely be identified and are not keyed out; it is important to distinguish septation with thickened cell walls, called distosepta, from simple eusepta; also it is important to observe whether or not the hamathecium filaments = paraphyses are branched; in case one of these characters is not ascertained, try both alternatives)

Please note that **keys A to E** mainly contain Pyrenulaceae, and **keys F to Ω** are all Trypetheliaceae. The keys look similar to previously published keys, but differ in numerous points, notably additional species and the corrections of mistakes. Each key alternative is given a separate code (like K2b) for easy future reference. Updated known world distributions are indicated for each species.

1a Hamathecium filaments absent or slimy; algae dark green, verrucarioid; thallus usually microsquamulose. Verrucariaceae (corticolous: *Agonimia*, *Endocarpon*, *Placidium*, *Psoroglaena*)

1b Hamathecium filaments present; algae trentepohlioid, or sometimes absent. 2

2a Ascospores simple. 3

2b Ascospores septate. 5

3a Ascospores globose, >50 per ascus; pantropical (Brazil, Papua New Guinea, Solomon Islands). *Melanophloea pacifica* P. James & Vězda

3b Ascospores 1 to 8 per ascus. 4

4a Ascospores ornamented with warts or spines. *Monoblastia*

4b Ascospores not ornamented, with wide, shaped gelatinous sheath: pantropical, but rarely reported (Brazil, Papua New Guinea). *Papilionovela athallina* Aptroot

5a Hamathecium with unbranched paraphyses or hamathecium unclear. 6

5b Hamathecium with branched to anastomosing paraphysoids. 19

6a Ascospores brown. 7

6b Ascospores hyaline. 12

7a Ascospores remaining stuck in mass on top of the ascoma, mazaedioid. Pyrenulaceae (*Mazaediothecium*, *Pyrgillus*). **Key E**

7b Ascospores discharged. 8

8a Ascospores 1-septate. 9

8b Ascospores more than 1-septate. Pyrenulaceae. 11

9a Ascospores not ornamented, often thick-walled, distoseptate. Pyrenulaceae (*Clypeopyrenis*, *Distopyrenis*, *Granulopyrenis*, *Parapyrenis*). **Key D**

9b Ascospores ornamented, thin-walled. 10

10a Ascospores up to $20 \times 8 \mu\text{m}$, only in one species larger (up to $30 \times 10 \mu\text{m}$). *Pseudobogoriella*. **Key X**

10b Ascospores $20\text{--}50 \times 8\text{--}15 \mu\text{m}$. *Bogoriella*. **Key Q**

11a Ascospores transversely septate. Pyrenulaceae (mostly *Eopyrenula*, *Lithothelium*, *Pyrenula*). **Key B**

11b Ascospores (sub)muriform. Pyrenulaceae (mostly *Anthracothecium*, *Pyrenula*). **Key A**

12a Ascospores >10 per ascus. *Thelopsis*

12b Ascospores 1 to 8 per ascus. 13

13a Hamathecium a pseudoparenchymatous tissue; ascomata usually in dense groups with common black cover. *Mycoporum*

13b Hamathecium consisting of filaments. 14

-
- 14a Ascospores thick-walled, distoseptate, either with rather unclear septation or with filiform ascospores). Pyrenulaceae (*Celothelium*, *Lithothelium*). **Key C**
- 14b Ascospores thin-walled, euseptate. 15
- 15a Ascomata in thalline warts with black pulveraceous mass. *Phyllobathelium*
- 15b Ascomata not in thalline warts with black pulveraceous mass. 16
- 16a Thallus medulla bright yellow or orange. *Myeloconis*
- 16b Thallus medulla not bright yellow or orange. 17
- 17a Ascomata with periphyses in addition to paraphyses. *Topelia*
- 17b Ascomata without periphyses. 18
- 18a Ascomata adorned with setae formed of conglutinated hairs. *Trichothelium*
- 18b Ascomata usually not adorned, but if with setae, these not made of conglutinated hairs. *Porina*
- 19a Ascospores distoseptate, mostly thick-walled; when relatively thin-walled, either hyaline and with constrictions or brown and ornamented. Trypetheliaceae. 23
- 19b Ascospores thin-walled, euseptate. 20
- 20a Hamathecium filaments unbranched between the asci. 21
- 20b Hamathecium filaments branched. 22
- 21a Ascospores 1 to 3-septate. Monoblastiaceae (*Acrocordia*, *Anisomeridium*, *Megalotremis*, *Trypetheliopsis*)
- 21b Ascospores muriform. *Julella*
- 22a Ascospores hyaline, usually uniseriate in the ascus; conidia with gelatinous appendages. Strigulaceae (*Dichoporis*, *Flagellostrigula*, *Strigula*, *Swinscowia*)
- 22b Ascospores biseriate; conidia without appendages. Thelenellaceae (*Aspidothecium*, *Thelenella*)
- 23a Ascospores remaining hyaline. 24
- 23b Ascospores becoming brown. 50
- 24a Ascospores 1-septate. 25
- 24b Ascospores 3-septate to muriform. 26
- 25a Ascospores smooth-walled. *Constrictolumina*. **Key R**
- 25b Ascospores granular ornamented. *Macroconstrictolumina*. **Key T**

- 26a Thallus ecorticate, usually whitish or greyish. 27
26b Thallus distinctly corticate, olive-green to yellowish brown. 31
- 27a Hamathecium filaments basally thickened, branched but usually not anastomosing.
28
27b Hamathecium filaments thin, straight, branched and anastomosing to form a net-like
structure. 29
- 28a Ascospores smooth-walled. *Constrictolumina*. **Key R**
28b Ascospores granular ornamented. *Macroconstrictolumina*. **Key T**
- 29a Ascospores with diamond-shaped lumina. *Pseudopyrenula*. **Key Y**
29b Ascospores with almost rectangular lumina. 30
- 30a Ascospores transversely septate, or when muriform, ostiole apical. *Polymeridium*.
Key W
30b Ascospores muriform; ostioles lateral. *Dictyomeridium*. **Key S**
- 31a Ascospores transversely septate. 32
31b Ascospores muriform. 41
- 32a Ascospores very large with few (3–5) septa, typically over $100 \times 30 \mu\text{m}$ and up to
 $190 \times 60 \mu\text{m}$; septa and walls somewhat thickened but lumina not astrothelioid.
Architrypethelium. **Key G**
32b Ascospores when 3–5-septate very rarely exceeding $100 \times 30 \mu\text{m}$ and if so,
ascospores distinctly astrothelioid, with diamond-shaped lumina. 33
- 33a Ascospores with diamond-shaped lumina. 34
33b Ascospores with ellipsoid to lentiform or almost rectangular lumina. 38
- 34a Ascomata strongly prominent to sessile, completely exposed, pure black, somewhat
egg-shaped. *Nigrovothelium*. **Key V**
34b Ascomata immersed to erumpent or aggregate in erumpent to sessile pseudostromata.
Astrothelium. 35
- 35a Ascomata each with a separate, consistently apical ostiole. 36
35b Ascomata either with a separate, eccentric to lateral ostiole or several ascomata with
a fused ostiole which is then positioned apically relative to the ascomatal cluster
but with individual channels originating eccentric or lateral from each individual
ascoma or chamber. 37

- 36a Thallus and/or ascomata with external or internal, yellow to red pigment(s) (usually K+ red to purple) and/or with lichexanthone (then UV+ yellow). **Key H**
- 36b Pigments and lichexanthone absent. **Key I**
- 37a Thallus and/or ascomata with external or internal, yellow to red pigment(s) (usually K+ red to purple) and/or with lichexanthone (then UV+ yellow). **Key J**
- 37b Pigments and lichexanthone absent. **Key K**
- 38a Ascomata aggregate in prominent to sessile, brownish black pseudostromata; ascospores with thin septa and walls. *Bathelium*. **Key P**
- 38b Ascomata solitary to pseudostromatic but pseudostromata not as above; ascospores with slightly thickened septa. 39
- 39a Ascomata immersed to erumpent or rarely indistinctly pseudostromatic; pigments mostly absent. 40
- 39b Ascomata aggregate in distinct, prominent to sessile pseudostromata; internal and/or external pigments usually present. *Trypethelium*. **Key Z**
- 40a Thallus UV-negative. *Viridothelium*. **Key □**
- 40b Thallus UV+ yellow; Brazil. *Exiliseptum ocellatum* (Müll. Arg.) R.C. Harris
- 41a Ascospores with ellipsoid to lentiform or almost rectangular lumina. 42
- 41b Ascospores with diamond-shaped lumina. 45
- 42a Ascomata single. *Architrypethelium submuriforme* Aptroot
- 42b Ascomata aggregate. 43
- 43a Ascomata aggregate in prominent to sessile, brownish black pseudostromata. *Bathelium*. **Key P**
- 43b Ascomata in erumpent, whitish pseudostromata. 44
- 44a Thallus UV-negative: Panama. *Viridothelium tricolor* Lücking, M.P. Nelsen & N. Salazar
- 44b Thallus UV+ yellow; Brazil. *Trypethelium muriforme* Aptroot & M.F. Souza
- 45a Ascomata in prominent to sessile warts covered by a thick layer of yellow-orange or red pigment. *Marcelaria*. **Key U**
- 45b Ascomata immersed to erumpent, rarely prominent, usually covered by thallus, rarely with a thin layer of pigment pruina. 46

- 46a Ascomata aggregate in irregular, blackish pseudostromata; ascospores about $50 \times 15 \mu\text{m}$; Thailand. *Architrypethelium murisporum* Luangsaph., Lumbsch & Sangvichien
- 46b Ascomata and ascospores not with the above combination of characters.
Astrothelium. 47
- 47a Ascomata each with a separate, consistently apical ostiole. 48
- 47b Ascomata either with a separate, eccentric to lateral ostiole or several ascomata with a fusoid ostiole which is then positioned apically relative to the ascomatal cluster but with individual channels originating eccentric or lateral from each individual ascoma or chamber. 49
- 48a Thallus and/or ascomata with external or internal, yellow to red pigment(s) (usually K+ red to purple) and/or with lichexanthone (then UV+ yellow). **Key L**
- 48b Pigments and lichexanthone absent. **Key M**
- 49a Thallus and/or ascomata with external or internal, yellow to red pigment(s) (usually K+ red to purple) and/or with lichexanthone (then UV+ yellow). **Key N**
- 49b Pigments and lichexanthone absent. **Key O**
- 50a Ascospores transversely septate. 51
- 50b Ascospores (sub-)muriform. 56
- 51a Ascospores with a basal euseptum and the basal cell pale, otherwise distoseptate and brown; neotropical. *Polypyrenula sexlocularis* (Müll. Arg.) D. Hawksw.
- 51a Ascospores septa all similar. 52
- 52a Ascospores large, usually over $100 \mu\text{m}$ long. 53
- 52b Ascospores small, under $50 \mu\text{m}$ long. 54
- 53a Ascospores 3-septate, $25\text{--}50 \mu\text{m}$ broad, about 3–4 times as long as broad.
Architrypethelium. **Key G**
- 53b Ascospores 11–15-septate, $25\text{--}30 \mu\text{m}$ broad, about 4–5 times as long as broad;
Colombia. *Astrothelium fuscosporum* Soto-Medino, Aptroot & Lücking
- 54a Ascospores drop-shaped, macrocephalic, with strongly thickened terminal walls and two narrow, rectangular lumina near the center; Azores. *Schummia angulata* (Aptroot & Schumm) Lücking, R. Miranda & Aptroot
- 54b Ascospores ellipsoid, not macrocephalic. 55
- 55a Ascospores up to $20 \times 8 \mu\text{m}$, only in one species larger (up to $30 \times 10 \mu\text{m}$).
Pseudobogoriella. **Key X**

55b Ascospores $20\text{--}50 \times 8\text{--}15 \mu\text{m}$. *Bogoriella*. **Key Q**

56a Ascospores small, under $50 \times 20 \mu\text{m}$, often with irregularly thickened endospore and halter-shaped lumina. *Bogoriella*. **Key Q**

56b Ascospores very large, $150\text{--}400 \times 50\text{--}140 \mu\text{m}$, with outer wall often breaking under pressure. *Aptrootia*. **Key F**

Key A. Mostly tropical corticolous pyrenocarpous lichens with simple paraphyses and brown, submuriform to muriform, distoseptate ascospores (mostly Pyrenulaceae: *Anthracothecium*, *Lithothelium* and *Pyrenula*)

A1a Young ascospores with eusepta only; mature ascospores with scarce endospore and edgy corners. *Anthracothecium*. 2

A1b Septation only or mostly distoseptate, with notably thick endospore layers, in particular in the corners, and causing the locules to appear rounded. 6

A2a Ascomata simple. 3

A2b Ascomata mostly compound with a joint ostiole. 5

A3a Ostiole apical. 4

A3b Ostiole lateral; pantropical. *Anthracothecium australiense* (Müll. Arg.) Aptroot

A4a Ascospores 2–4/ascus; pantropical. *Anthracothecium macrosporum* (Hepp) Müll. Arg.

A4b Ascospores 6–8/ascus; pantropical. *Anthracothecium prasinum* (Eschw.) R.C. Harris

A5a Ascospores 1–2/ascus; pantropical. *Anthracothecium interlatens* (Nyl.) Aptroot

A5b Ascospores 6–8/ascus; australasian, possibly also african.

Anthracothecium gregale (C. Knight) Aptroot

A6a Ascospores strongly flattened (lozenge-shaped), with two rows of locules.
Sulcopyrenula. 7

A6b Ascospores not flattened, circular in transverse section. 11

A7a Thallus UV-negative. 8

A7b Thallus UV+ yellow. 9

A8a Hamathecium interspersed; neotropical. *Sulcopyrenula canellae-albae* (Fée) H. Harada

A8b Hamathecium not interspersed; pantropical. *Sulcopyrenula staurospora* (Tuck.) H. Harada

- A9a Ascospores with 4 locules. 10
A9b Ascospores with 8 locules; Guyana. *Sulcopyrenula biseriata* Aptroot & Sipman
- A10a Ascospores ellipsoidal, c. 2 times as long as wide; Brazil.
Sulcopyrenula cruciata Aptroot
A10b Ascospores nearly globose; neotropical. *Sulcopyrenula subglobosa* (Riddle)
Aptroot
- A11a Ascospores submuriform, only one or a few of the median locules with longitudinal septa. 12
A11b Ascospores muriform (when ascospores small, logically only few longitudinal septa can be present; when in doubt, start with the first alternative). 25
- A12a Ascomata usually fused or compound, when separate, ostioles lateral; ostioles apical or lateral. 13
A12b Ascomata separate and simple; ostioles apical. 18
- A13a Ascomata fused, in black, exposed groups with several ostioles; ascospores 15–22 μm long; australasian. *Lithothelium hieroglyphicum* Aptroot
A13b Ascomata compound, with several, usually flask-shaped chambers usually immersed in the thallus and with a joint ostiole. 14
- A14a Ascospores mostly less than 25 μm long. 15
A14b Ascospores mostly over 25 μm long. 17
- A15a Thallus with pseudocyphellae; ascospores 15–19 μm long; Guyana. *Pyrenula infrastroidea* Aptroot & Sipman
A15b Thallus without pseudocyphellae. 16
- A16a Ascospores red-brown, 14–23 μm long; eastern palaeotropical (Australia to Japan).
Lithothelium nanosporum (C. Knight) Aptroot
A16b Ascospores grey-brown, 17–20(–25) μm long; neotropical. *Pyrenula subvariabilis*
Aptroot & Sipman
- A17a Ascospores up to 32 μm long; australasian. *Pyrenula subumbilicata* (C. Knight)
Aptroot
A17b Ascospores 45–65 μm long; neotropical. *Pyrenula erumpens* R.C. Harris
- A18a Ascospores 13–35(–40) μm long. 19
A18b Ascospores 34–72 μm long. 24
- A19a Ascospores 13–17 μm long. 20

- A19b Ascospores 22–35(-40) μm long. 21
- A20a Ascospores with little endospore formation, lumina almost angular; on wood, pacific. *Pyrenographa irregularis* (Wehmeyer) R.C. Harris
- A20b Ascospores much thickened, with rounded lumina; Florida. *Distopyrenis submuriformis* R.C. Harris
- A21a Thallus probably without algae; temperate. *Requienella seminuda* (Pers.) Boise
- A21b Thallus superficial, with copious algae. 22
- A22a Ascospores 22–40 μm long; overmature ascospores with reddish content before shiveling; pantropical. *Pyrenula seminuda* (Müll. Arg.) Sipman & Aptroot
- A22b Ascospores 23–35 μm long; overmature spores all shrivelled. 23
- A23a Ascospores 23–35 μm long; hamathecium not inspersed; eastern palaeotropical. *Pyrenula gibberulosa* (Vain.) Aptroot
- A23b Ascospores 23–33 μm long; hamathecium inspersed; eastern palaeotropical (India). *Pyrenula darjeelingensis* Jagadeesh Ram & G.P. Sinha
- A24a Ascospores with pointed ends; hamathecium not inspersed; eastern palaeotropical (Australia to Japan). *Pyrenula subvariolosa* (C. Knight) Aptroot
- A24b Ascospores with rounded ends; hamathecium inspersed; neotropical. *Pyrenula novemseptata* Vain.
- A25a Ascomata and/or thallus or medulla with yellow, orange or red anthraquinone pigments; pigments K+ pink to purplish. 26
- A25b Anthraquinones absent; thallus K– or yellowish. 32
- A26a Thallus warts internally with soft orange medulla; Philippines. *Pyrenula endocrocea* Aptroot
- A26b Yellow, orange or red pigment on outside. 27
- A27a Ascomata and/or thallus red. 28
- A27b Ascomata and/or thallus yellow to orange. 30
- A28a Ostiole apical; hamathecium inspersed; neotropical. *Pyrenula cruentata* (Müll. Arg.) R.C. Harris
- A28b Ostiole lateral; hamathecium not inspersed. 29
- A29a Ascomata partly fused with joint ostioles; ascospores with up to 6 locules per segment; caribbean. *Pyrenula kermesina* R.C. Harris

- A29b Ascomata solitary; ascospores with up to 2 locules per segment; pacific. *Pyrenula palmarum* (Krempelh.) R.C. Harris
- A30a Ascospores 10–23 µm long, with 3 primary septa. 31
A30b Ascospores 23–35 µm long, with 5–7 primary septa; neotropical.
Pyrenula ochraceoflavens (Nyl.) R.C. Harris
- A31a Thallus red-orange; ascospores 9–18 µm long; Brazil. *Pyrenula aurantiacorubra*
Aptroot & M. Cáceres
A31b Thallus yellow to orange; ascospores 10–23 µm long; pantropical.
Pyrenula ochraceoflava (Nyl.) R.C. Harris
(with small ascospores with two layers of locules; pacific: *Pyrenula ochraceoflava* var.
pacifica P.M. McCarthy)
- A32a Ostioles lateral. 33
A32b Ostioles apical. 41
- A33a Ascospores <70 µm long. 34
A33b Ascospores >70 µm long. 36
- A34a Ascomata at least partly with more than one chamber and with joint ostiole. 35
A34b Ascomata all simple and single, with separate ostioles; ascospores 15–33 µm long;
pantropical (El Salvador, India). *Pyrenula microspora* (Nagarkar & Patw.) Upreti
- A35a Ascospores 25–45 µm long; pantropical. *Pyrenula astroidea* (Fée) R.C. Harris
A35b Ascospores 45–70 µm long; pantropical. *Pyrenula ravenelii* (Tuck.) R.C. Harris
- A36a Ascospores 2/ascus, 135–200 µm long; pantropical. *Pyrenula lyoni* (Zahlbr.)
Aptroot
A36b Ascospores 4–8/ascus. 37
- A37a Ascospores 25–110 µm long. 38
A37b Ascospores 85–135 µm long. 40
- A38a Ascospores 25–31 µm long; Guyana. *Pyrenula triangularis* Aptroot & Sipman
A38b Ascospores 50–110(–135) µm long. 39
- A39a Ascospores 50–90 µm long; pantropical (El Salvador, India). *Pyrenula ceylonensis*
(Ajay Singh & Upreti) Aptroot
A39b Ascospores 70–135 µm long; W. Europe, Madagascar & Macaronesia). *Pyrenula hibernica* (Nyl.) Aptroot

-
- A40a Ascomata mostly with several chambers; pantropical. *Pyrenula schiffneri* (Zahlbr.) Aptroot
- A40b Ascomata often with single chamber; neotropical. *Pyrenula chilensis* (Fée) R.C. Harris
- A41a Ascospores <25 µm long. 42
- A41b Ascospores >25 µm long. 46
- A42a Hamathecium inspersed. 43
- A42b Hamathecium not inspersed. 44
- A43a Ascospores 13–18 µm long, with basal cilium; Mauritius. *Pyrenula muriciliata* Diederich & Ertz
- A43b Ascospores 20–25 µm long, without cilium; Borneo. *Pyrenula borneensis* Aptroot
- A44a Thallus UV+yellow; pantropical. *Pyrenula confinis* (Nyl.) R.C. Harris
- A44b Thallus UV-negative. 45
- A45a Ascospores 15–22 µm long; pantropical. *Pyrenula parvinuclea* (Meyen & Flot.) Aptroot
- A45b Ascospores 7–10 µm long; India. *Pyrenula nanospora* (Ajay Singh) Upreti
- A46a Old ascospores containing orange oil before shriveling; pantropical. *Pyrenula breutelii* (Müll. Arg.) Aptroot
- A46b Old ascospores without orange oil, directly shriveling. 47
- A47a Ascospores >80 µm long, mostly 2/ascus. 48
- A47b Ascospores <80 µm long, mostly 4–8/ascus. 53
- A48a Hamathecium inspersed, ascospores 90–200 µm long. 49
- A48b Hamathecium not inspersed. 50
- A49a Thallus UV+ yellow, with lichexanthone; neotropical. *Pyrenula xanthoglobulifera* Aptroot, Lücking & M. Cáceres
- A49b Thallus UV-negative, without lichexanthone; pantropical. *Pyrenula globifera* (Eschw.) Aptroot
- A50a Thallus without pseudocyphellae, ascospores 80–140(–155) µm long; pantropical. *Pyrenula platystoma* (Müll. Arg.) Aptroot
- A50b Thallus with pseudocyphellae. 51
- A51a Ascospores 80–180 µm long, 2-8/ascus. 52

- A51b Ascospores 205–350 µm long, 1/ascus; Guyana. *Pyrenula monospora* Aptroot & Sipman
- A52a Ascospores 80–110 µm long, 2-8/ascus; Hawaii. *Pyrenula neosandwicensis* Aptroot
- A52b Ascospores 115–180 µm long, 2/ascus; pantropical. *Pyrenula duplicans* (Nyl.) Aptroot
- A53a Hamathecium inspersed. 54
- A53b Hamathecium not inspersed. 55
- A54a Ascospores 2/ascus, 55–75 µm long; Brazil. *Pyrenula bispora* Aptroot & M. Cáceres
- A54b Ascospores 8/ascus, 30–45 µm long; pantropical (India, Brazil). *Pyrenula sublaevigata* (Patw. & Makhija) Upreti
- A55a Locules relatively large and angular, with up to 6 between 2 primary septa; pantropical. *Pyrenula leucostoma* Ach.
- A55b. Locules mostly round, at least in the central part of the ascospore with more than 6 between 2 primary septa. 56
- A56a Ascospores <50 µm long. 57
- A56b Ascospores >50 µm long. 60
- A57a Ascospores 25–35 µm long; palaeotropical. *Pyrenula welwitschii* (Upreti & Ajay Singh) Aptroot
- A57b Ascospores >35 µm long. 58
- A58a Ascospores 11–15 µm wide; pantropical. *Pyrenula thelomorpha* Tuck.
- A58b Ascomata 14–22 µm wide. 59
- A59a Old ascospores containing colourless oil before shriveling; neotropical. *Pyrenula oleosa* R.C. Harris
- A59b Old ascospores without oil, directly shriveling when old; neotropical. *Pyrenula dissimulans* (Müll. Arg.) R.C. Harris
- A60a Ascomata deeply immersed in bark below the thallus, not visible, only ostioles visible; Brazil. *Pyrenula abditicarpa* Aptroot & M. Cáceres
- A60b Ascomata more-or-less exposed. 61
- A61a Ascospores with rounded ends; pantropical. *Pyrenula pyrenuloides* (Mont.) R.C. Harris

A61b Ascospores with pointed ends; pantropical. *Pyrenula papillifera* (Nyl.) Aptroot

Key B. Corticolous pyrenocarpous lichens with simple paraphyses and brown, transversely 3- or more septate distoseptate ascospores (mostly Pyrenulaceae: *Eopyrenula*, *Lithothelium* and *Pyrenula*)

B1a Ascospores 3-loculate. 2

B1b Ascospores 4- or more-loculate. 4

B2a Ascospore septation strongly asymmetrical; Madagascar. *Lacrymospora parasitica*
Aptroot

B2b Ascospore septation symmetrical. 3

B3a Ascomata aggregated in grey to black pseudostromata; pantropical. *Pyrenula lineatostroma* Aptroot

B3b Ascomata dispersed, not stromatoid; Brazil. *Pyrenula biseptata* Aptroot & M. Cáceres

B4a Ascospores nearly only euseptate, locules rectangular, at most a bit rounded in the corners, end cells often paler than middle cells. 5

B4b Ascospores clearly distoseptate, locules more or less rounded or diamond-shaped. 17

B5a Ascospores more than 5 times as long as wide; South Africa. *Pyrenowilmsia wilmsiana* (Zahlbr.) Aptroot

B5b Ascospores less than 4 times as long as wide. 6

B6a Ascospores only 4-loculate; macroconidia 3-septate or unknown. 7

B6b Ascospores 4–8-loculate; macroconidia either 1–7-septate or unknown. 12

B7a Ascoma wall dense. 8

B7a Ascoma wall cellular. *Eopyrenula*. 10

B8a Ascospores smooth, 14–17 µm long; neotropical. *Pyrenula tenuisepta* R.C. Harris

B8b Ascospores verrucose, 30–47 µm long. *Bogoriella*. 9

B9a Ascomata at least partly fused with a more or less joint ostiole; Japan; pantropical. *Bogoriella collospora* (Vain.) Aptroot & Lücking

B9b Ascomata all single; Australasian; pantropical. *Bogoriella queenslandica* (Müll. Arg.) Aptroot & Lücking

- B10a Ascospores 4–5 µm wide; New world temperate. *Eopyrenula parvispora* R.C. Harris & Aptroot
B10b Ascospores 5.5–7.5 µm wide. 11
- B11a Ascospores mostly <15 µm long; Atlantic (W. Europe). *Eopyrenula avellanae* Coppins
B11b Ascospores mostly >15 µm long; Atlantic (W. Europe). *Eopyrenula grandicula* Coppins
- B12a Ascospores 4(–6)-loculate, macroconidia 1-septate; Old world temperate. *Eopyrenula leucoplaca* (Wallr.) R.C. Harris
B12b Ascospores (4–)6–8-loculate; macroconidia 3–7-septate or unknown. 13
- B13a Ascospores mostly >15 µm long. *Eopyrenula*. 14
B13b Ascospores mostly over 25 µm long. 15
- B14a Ascospores >6 µm wide, macroconidia (5–)7-septate; Atlantic Europe. *Eopyrenula septemseptata* Coppins
B14b Ascospores <6 µm wide, macroconidia 3(–4)-septate; New world temperate. *Eopyrenula intermedia* Coppins
- B15a Ascospores mostly 37–47 µm long, constricted at the septa; Europe. *Blastodesmia nitida* A. Massal.
B15b Ascospores 25–35 µm long. *Requienella*. 16
- B16a Ascospores 5-7-septate, rarely 3-septate; temperate. *Requienella seminuda* (Pers.) Boise
B16b Ascospores mostly 3-septate; temperate. *Requienella fraxini* Jaklitsch & Voglmayr
- B17a Ascospores red brown and locules at least becoming rounded when older (species on rock not keyed out here). *Lithothelium*. 18
B17b Ascospores grey to brown, rarely red brown and then locules angular or diamond-shaped. *Pyrenula*. 22
- B18a Thallus UV+ yellow; Costa Rica. *Lithothelium fluorescens* Aptroot & Sipman
B18b Thallus UV-negative. 19
- B19a Ascospores 4-loculate. 20
B19b Ascospores 8-loculate. 21
- B20a Ascospores 15–20 µm long; eastern palaeotropical. *Lithothelium decumbens* (Müll. Arg.) Aptroot

-
- B20b Ascospores 25–40 µm long; northern temperate. *Lithothelium phaeosporum* (R.C. Harris) Aptroot
- B21a Ascospores 30–40 µm long; northern temperate. *Lithothelium septemseptatum* (R.C. Harris) Aptroot
- B21b Ascospores 55–80 µm long; New world temperate. *Lithothelium macrosporum* (R.C. Harris) Aptroot
- B22a Ostioles pointing in various directions, mostly eccentric to lateral; ascomata sometimes with several chambers connected to joint ostioles. 23
- B22b Ostioles apical or, when eccentric, all pointing in the same direction; ascomata with one chamber or each chamber with own ostiole. 48
- B23a Ascospores 6–16-loculate. 24
- B23b Ascospores 4-loculate. 27
- B24a Ascospores 6-loculate. 25
- B24b Ascospores 10–16-loculate. 26
- B25a Ascospores 42–55 µm long, 8/ascus; neotropical (Colombia, Florida). *Pyrenula pleiomeria* (Nyl.) Zahlbr.
- B25b Ascospores 60–65 µm long, 4/ascus; Venezuela. *Pyrenula tetraspora* Aptroot & Sipman
- B26a Ascospores 10–14-loculate, 65–90 × 17–22 µm; ascomata with single chamber; Brazil. *Pyrenula fusispora* (Malme) Aptroot
- B26b Ascospores 12–16-loculate, 50–70 × 4.5–6 µm; ascomata with 5–15 stellately fused chambers; Japan. *Pyrenula tokyensis* (Müll. Arg.) H. Harada
- B27a Thallus UV+ yellow. 28
- B27b Thallus UV-negative. 29
- B28a Ascospores 21–23 µm long; Hawaii. *Pyrenula hawaiiensis* Aptroot
- B28b Ascospores 32–45 µm long; Brazil. *Pyrenula crassiuscula* (Malme) Aptroot
- B29a Terminal locules directly against the exospore wall. 30
- B29b Terminal locules separated from the exospore wall by endospore thickening. 34
- B30a Ascomata at least partly with 2-7 chambers with joint ostioles; neotropical. *Pyrenula subgregantula* Müll. Arg.
- B30b Ascomata single. 31

- B31a Ascospores 13–25 µm long. 32
B31b Ascospores 25–65 µm long. 33
- B32a Ascospores 16–25 µm long; pantropical. *Pyrenula circumfiniens* Vain.
B32b Ascospores 13–16 µm long; neotropical. *Pyrenula elliptica* Müll. Arg.
- B33a Ascospores 45–65 µm long; neotropical. *Pyrenula erumpens* R.C. Harris
B33b Ascospores 25–35 µm long; pantropical. *Pyrenula cuyabensis* (Malme) R.C. Harris
- B34a Ostiole with red colour (inconspicuous character!). 35
B34b Ostiole without red colour. 36
- B35a Ostiole outside with a red ring; New world temperate. *Pyrenula wetmorei* R.C. Harris
B35b Ostiole red brown to bright red, eccentric, single; Brazil. *Pyrenula rubrolateralis* Aptroot & M. Cáceres
B35c Ostiole red, fused; Brazil. *Pyrenula sanguineoastroidea* Aptroot
- B36a Ascomata mostly with several chambers and a joint ostiole. 37
B36b Ascomata nearly all single. 43
- B37a Ascospores 10–24 µm long. 38
B37b Ascospores >25 µm long. 41
- B38a Ascospores 10–12 µm long; Argentina. *Pyrenula punctoleucotrypa* Aptroot, L.I. Ferraro & M. Cáceres
B38b Ascospores mostly 15–20 µm long. 39
- B39a Ascospores 17–20 x 5.5–6.5 µm; hamathecium interspersed with colourless oil globules instead of clear; Brazil. *Pyrenula inspersicollaris* Aptroot & M. Cáceres
B39b Ascospores 12–24 µm long; pantropical. 40
- B40a Thallus partly wine-red to purplish by K⁺ purple pigment; ascospores 16–21 µm long; Brazil. *Pyrenula violaceoastroidea* C.O. Mendonça, Aptroot & M. Cáceres
B40b Thallus not pigmented; ascospores 12–24 µm long; pantropical. *Pyrenula septicollaris* (Eschw.) R.C. Harris
- B41a Ascoma chambers deeply immersed; ascospores 23–27 µm long; Brazil. *Pyrenula leptaleoides* Aptroot & M. Cáceres
B41b Ascoma chambers emergent; ascospores mostly over 25 µm long. 42
- B42a Ascospores 25–35 µm long; pantropical. *Pyrenula cubana* (Müll. Arg.) R.C. Harris

-
- B42b Ascospores 30–45 μm long; pantropical. *Pyrenula cryptothelia* (Müll. Arg.) Aptroot & Etayo
- B43a Ascospores 13–17 μm long; hamathecium inspersed; USA (Florida). *Pyrenula wheeleri* R.C. Harris
- B43b Ascospores >17 μm long; hamathecium not inspersed. 44
- B44a Ascospores 17–28 μm long. 45
- B44b Ascospores 28–45 μm long (compare also *Pyrenula minarum*, with partly eccentric ostioles). 47
- B45a Ascospores with dark bands of euseptum material between the locules, 19–22 μm long; Brazil. *Pyrenula viridipyrgilla* Aptroot & M. Cáceres
- B45b Ascospores without dark bands, 18–30 μm long. 46
- B46a Ascospores with at least one pointed end; almost cosmopolitan. *Pyrenula acutispora* Kalb & Hafellner
- B46a Ascospores with rounded ends; pantropical. *Pyrenula microtheca* R.C. Harris
- B47a Ascospores 35–45 μm long; pantropical. *Pyrenula adacta* Fée
- B47b Ascospores 28–32 μm long; probably pantropical (only known from Brazil and Papua New Guinea). *Pyrenula gahavisukana* Aptroot
- B48a Ascospores with 2-8 or many hyaline cilia at the ends (not to be confused with germ tubes which are one per lumen). 49
- B48b Ascospores without cilia. 50
- B49a Ascospores 4-loculate; Australasian (Papua New Guinea). *Pyrenula ciliata* Aptroot
- B49b Ascospores 6–8-loculate; neotropical. *Pyrenula hirsuta* Etayo
- B50a Ascospores at least partly with more than 3 septa, or with 3 septa and long tails at one or both ends, seemingly more than 4-loculate. 51
- B50b Ascospores all distinctly 4-loculate. 65
- B51a Ascospores at least seemingly 5–8-loculate. 52
- B51b Ascospores 8–18-loculate, more than 4 times as long as wide. 59
- B52a Ascospore with tails c. 1 μm wide and not septate; palaeotropical. 53
- B52b Ascospore without tails, at most pointed; neotropical. 54
- B53a Ascospores with tails at both ends; Australia. *Pyrenula bicuspidata* Müll. Arg.
- B53b Ascospores with tail at basal end only; Oceania. *Pyrenula mastigophora* Aptroot

- B54a Ascospores more than 4 times as long as wide; neotropical. 55
B54b Ascospores less than 4 times as long as wide. 56
- B55a Ascospores more than 4 times as long as wide; hamathecium clear. *Pyrenula melanophthalma* (Mont.) Trevis.
B55b Ascospores 30–37 x 3–4 µm; thallus UV+ yellow instead of UV–; hamathecium interspersed with colourless oil globules. *Pyrenula musaespora* Aptroot & M. Cáceres
- B56a Ascospores <30 µm long. 57
B56b Ascospores >29 µm long. 58
- B57a Old ascospores with orange oil; thallus often with pseudocyphellae; pantropical. *Pyrenula sexlocularis* (Nyl.) Müll. Arg.
B57b Ascospores without orange oil; thallus without pseudocyphellae; South Africa. *Pyrenula wilmsiana* Müll. Arg.
- B58a Ascospores 29–35 µm long; hamathecium not interspersed; Australasian (Papua New Guinea). *Pyrenula sexluminata* Aptroot
B58b Ascospores 30–55 µm long; hamathecium interspersed; pantropical. *Pyrenula caracasana* Müll. Arg.
- B59a Ascospores with a long tail at one end; Pacific (Mariana Islands). *Pyrenula flagellata* H. Harada
B59b Ascospores without tails. 60
- B60a Ascospores 8–18-loculate. 61
B60b Ascospores (6–)8-loculate. 64
- B61a Thallus without pseudocyphellae. 62
B61b Thallus with pseudocyphellae. 63
- B62a Ascospores 13–18-loculate; eastern palaeotropical (Ryukyu Islands, Japan). *Pyrenula cylindrica* Kashiw.
B62b Ascospores 8–13-loculate; USA (Florida). *Pyrenula clavatispora* Common & Aptroot
- B63a Ascospores 8–12-loculate; eastern palaeotropical (India). *Pyrenula subcylindrica* Jagadeesh Ram & Upreti
B63b Ascospores 12–18-loculate, 75–90 × 4.5–6.5 µm, gradually attenuated toward the ends; New Caledonia. *Pyrenula filiformis* Aptroot

- B64a Ascospores 30–38 μm long; pantropical. *Pyrenula montagnei* Müll. Arg.
B64b Ascospores 53–70 μm long; Australasian (New Zealand), also neotropical?
Pyrenula moniliformis (C. Knight) Müll. Arg.
- B65a Ascospores more than 4 times as long as wide; Brazil. *Pyrenula fusoluminata*
Aptroot
B65b Ascospores less than 4 times as long as wide. 66
- B66a Ascospores mostly >50 μm long. 67
B66b Ascospores mostly <50 μm long. 73
- B67a Locules not in a straight line, somewhat zig-zag; Brazil. *Pyrenula hoehneliana*
Zahlbr.
B67b Locules in a straight line or ascospores bent, not zig-zag. 68
- B68a Terminal locules directly against the exospore wall. 69
B68b Terminal locules separated from the exospore wall by endospore thickening; thallus
with pseudocyphellae or crystalline papillae. 71
- B69a Thallus UV+yellow; neotropical. *Pyrenula praelucida* (Mont.) Trevis.
B69b Thallus UV-negative. 70
- B70a Hamathecium not inspersed; pantropical. *Pyrenula rockii* Zahlbr.
B70b Hamathecium inspersed; neotropical. *Pyrenula subpraelucida* Müll. Arg.
- B71a Hamathecium inspersed with orange oil globules, K+ greenish, not clear;
ascospores 60–75 x 24–28 μm ; thallus without papillae; Venezuela. *Pyrenula*
aurantioinspersa Aptroot & Sipman
B71b Hamathecium not inspersed. 72
- B72a Ascospores (45–)50–60 μm long; thallus without papillae but with
pseudocyphellae; pantropical. *Pyrenula immissa* (Stirt.) Zahlbr.
B72b Ascospores 75–90 μm long; thallus with crystalline papillae; Costa Rica. *Pyrenula*
montocensis Lücking
- B73a Locules not in a straight line, somewhat to strongly zig-zag. 74
B73b Locules in a straight line, not zig-zag. 75
- B74a Thallus UV+ yellow; pantropical. *Pyrenula aggregans* Vain.
B74b Thallus UV-negative; pantropical. *Pyrenula papilligera* (Leight.) Müll. Arg.

-
- B75a Ascomata and/or thallus with yellow, orange or red anthraquinones on the outside. 76
B75b Ascomata and thallus without external pigments. 97
- B76a Terminal locules directly against the exospore wall. 77
B76b Terminal locules separated from the exospore wall by endospore thickening. 80
- B77a Ascomata at least partly yellow; hamathecium not inspersed; neotropical and
pacific (Panama, Costa Rica, Puerto Rico, Hawaii). *Pyrenula luteopruinosa* Etayo
& Aptroot
B77b Ascomata or thallus orange or red; hamathecium inspersed. 78
- B78a Ascomata red; thallus UV-negative; Australasian (Lord Howe Island). *Pyrenula
howeana* Aptroot
B78b Thallus or ascomata orange, thallus UV+ yellow. 79
- B79a Ascomata orange; ascospores 17–20 µm long; eastern palaeotropical (Korea).
Pyrenula sipmanii Aptroot & K.H. Moon
B79b Thallus orange; ascospores 19–22 µm long; eastern palaeotropical (Thailand).
Pyrenula aurantiopileata Aptroot
- B80a Thallus yellow (in shade) to orange (in full light). 81
B80b Thallus and/or ascomata at least partly pink, red or lilac. 82
- B81a Ascospores ca. 25-40 µm long, not plicate; found in mangrove; neotropical.
Pyrenula cerina Eschw.
B81b Ascospores 55-65 µm long, longitudinally plicate; Chile. *Pyrenula plicata* Sipman
& Aptroot
- B82a Thallus and/or ascomata at least partly red. 83
B82b Thallus lilac; Brazil. *Pyrenula lilacina* C.O. Mendonça, Aptroot & M. Cáceres
B82c Thallus salmon pink; hamathecium inspersed; ascospores 24–27 × 13–16 µm;
Brazil. *Pyrenula salmonea* Aptroot
- B83a Hamathecium not inspersed. 84
B83b Hamathecium inspersed with colourless oil globules. 91
- B84a Ascomata aggregated in pseudostromata. 85
B84b Ascomata mostly separate. 88
- B85a Ascospores 15–18(-20) µm long. 86
B85b Ascospores 20–24 µm long; ascomata densely aggregated in well-delimited, bright
red pseudostromata; Brazil. *Pyrenula sanguinea* Aptroot, M. Cáceres & Lücking

B86a Ascomata covered by red thallus; Brazil. *Pyrenula cinnabarina* Aptroot, E.L. Lima & M. Cáceres

B86b Ascomata covered by red pigment only. 87

B87a Ascospores 15–17 μm long; Costa Rica. *Pyrenula rubroanomala* Aptroot & Lücking

B87b Ascospores 15–18(–20) μm long; Brazil. *Pyrenula coccinea* C.O. Mendonça, Aptroot & M. Cáceres

B88a Ascospores 16–19 μm long. 89

B88b Ascospores 21–28 μm long. 90

B89a Red colouration only in tissue around the ostiole; neotropical. *Pyrenula rubrostoma* R.C. Harris

B89b Red coloration extending over the thallus; Brazil. *Pyrenula rubronitidula* Aptroot & M. Cáceres

B90a Thallus extensively carmine-red-stained; Brazil. *Pyrenula rubroacutispora* Aptroot

B90b Only ostioles of (immersed) ascomata red; Caribbean. *Pyrenula sanguineostiolata* Aptroot & Mercado Diaz

B91a Ascomata aggregated in pseudostromata; ascospores 12–15 x 6–8 μm ; Brazil. *Pyrenula reginae* E.L. Lima, Aptroot & M. Cáceres

B91b Ascomata not aggregated. 92

B92a Ascospores about 15–18 μm long. 93

B92b Ascospores mostly over 19 μm long. 94

B93a Ascospores about 19–22 μm long. 95

B93b Ascospores about 19–22 μm long. 96

B94a Thallus orange-brown to red, ostioles black; Brazil. *Pyrenula rubromamillana* E.L. Lima, Aptroot & M. Cáceres

B94b Only ostioles pigmented, red; Brazil. *Pyrenula rubrostigma* Aptroot & M. Cáceres

B95a Thallus dark reddish to orange-brown, K+ crimson; ascomata hemispherical, 0.4–0.7 mm diam.; eastern palaeotropical (Java). *Pyrenula rubrojavonica* Aptroot

B95b Thallus orange, K+ purple; ascomata low-conical, 0.7–1 mm diam.; Brazil. *Pyrenula aurantiacothallina* C.O. Mendonça, Aptroot & M. Cáceres

- B96a Ascospores 25–29 x 10–12 µm; thallus with red spots or lines; ascomata immersed in thallus, 0.2–0.4 mm diam.; Caribbean. *Pyrenula sanguineomeandrata* Aptroot & Mercado Diaz
- B96b Ascospores 27–35 µm long; thallus with red patches or completely red; ascomata erumpent, c. 0.4–0.8 mm diam.; pantropical. *Pyrenula cruenta* (Mont.) Vain.
- B97a Ascomata mostly aggregated, with fused walls but with separate ostioles. 98
- B97b Ascomata mostly simple, only aggregated as by chance when crowded (if difficult to decide, start with the first alternative). 112
- B98a Old ascospores with red oil; Puerto Rico. *Pyrenula concastroma* R.C. Harris
- B98b Old ascospores without red oil. 99
- B99a Terminal locules directly against the exospore wall; Australasian (Papua New Guinea). *Pyrenula pyrenastrospora* Aptroot
- B99b Terminal locules separated from the exospore wall by endospore thickening. 100
- B100a Hamathecium interspersed with colourless oil globules. 101
- B100b Hamathecium not interspersed. 103
- B101a Ascospores 16–20 µm long. 102
- B101b Ascospores 25–40 µm long; ostioles sometimes partly eccentric; pantropical. *Pyrenula minarum* Vain.
- B102a Ascomata hemispherical, 0.3–0.5 mm diam.; ascospores 16–18 µm long; Brazil. *Pyrenula paraminarum* Aptroot & M. Cáceres
- B102b Ascomata conical with flattened tip, 0.4–1.2 mm diam.; ascospores 17–20 µm long; Argentina. *Pyrenula inspersionotrypa* Aptroot, L.I. Ferraro & M. Cáceres
- B103a Ascospores mostly <20 µm long. 104
- B103b Ascospores mostly >21 µm long. 108
- B104a Ascospores with dark bands between the locules; USA (Florida). *Pyrenula atrolaminata* R.C. Harris
- B104b Ascospores without dark bands. 105
- B105a Ascomata not in dense stromata but more loosely aggregated, the individual ascomata often well recognizable, conical, with distinct ostioles. 106
- B105b Ascomata in dense, well delimited stromata, subglobose with indistinct, black ostioles. 107

- B106a Ascospores 11–13 x 3.5–5 μm ; Brazil. *Pyrenula aggregataspistea* Aptroot & M. Cáceres
- B106b Ascospores 15–18(–22) μm long; Brazil. *Pyrenula infraleucotrypa* Aptroot & M. Cáceres
- B107a Ascospores mostly >15 μm long; ascomata in dense stromata; pantropical. *Pyrenula anomala* (Ach.) Vain.
- B107b Ascospores mostly <15 μm long; pantropical. *Pyrenula arthoniotheca* Upreti
- B108a Ascospores mostly <25 μm long. 109
- B108b Ascospores mostly >25 μm long. 110
- B109a End locules elongated; Costa Rica. *Pyrenula subsoluta* (Müll. Arg.) Aptroot
- B109b All locules more or less rounded; eastern palaeotropical. *Pyrenula leucotrypa* (Nyl.) Upreti
- B110a Thallus with pseudocyphellae; neotropica. 1111
- B110b Thallus without pseudocyphellae; eastern palaeotropical (Sri Lanka). *Pyrenula zeylanica* Upreti & Ajay Singh
- B111a Thallus with pseudocyphellae; Cuba. *Pyrenula wrightii* (Müll. Arg.) R.C. Harris
- B111b Thallus with pseudocyphellae, ascomata slightly fused, in groups of few ascomata which are hardly touching each other; Brazil. *Pyrenula diamantinensis* C.O. Mendonça, Aptroot & M. Cáceres
- B112a Thallus ecorticate, whitish. 113
- B112b Thallus corticate (may be difficult to observe in poor specimens; when in doubt try the first alternative first). 118
- B113a Thallus UV+yellow. 114
- B113b Thallus UV–. 115
- B114a Ascospores 18–21 μm long; pantropical. *Pyrenula cocoes* Müll. Arg.
- B114b Ascospores 14–17 μm long; Brazil. *Pyrenula xanthinspersa* Aptroot & M. Cáceres
- B115a Hamathecium inspersed; pantropical (Brazil, Philippines). *Pyrenula albothallina* Vain.
- B115b Hamathecium not inspersed. 116
- B116a Thallus white to grey, dull; ascomata <0.7 mm diam.; pantropical. *Pyrenula microcarpa* Müll. Arg.

- B116b Thallus cream to silvery or inapparent, smooth; temperate. 117
- B117a Ascospores <18 µm long; hamathecium K-negative; pycnidia present. Known from temperate regions on the northern hemisphere and reported from Chile, so probably cosmopolitan in temperate regions. *Pyrenula coryli* A. Massal.
- B117b Ascospores >18 µm long; hamathecium sometimes KOH+ blue (fleeting), pycnidia absent; temperate northern hemisphere. *Pyrenula laevigata* (Pers.) Arnold
- B118a Old ascospores with orange oil. 119
- B118b Old ascospores without orange oil. 120
- B119a Ascospores 25–35 µm long; pantropical. *Pyrenula bahiana* Malme
- B119b Ascospores (30–)35–51 µm long; eastern palaeotropical. *Pyrenula thailandica* Aptroot
- B120a Terminal locules all directly against the exospore wall (if anthraquinones present on thallus and/or ascomata, compare B75). 121
- B120b Terminal locules mostly (at least in mature ascospores) separated from the exospore wall by endospore thickening. 142
- (See also *P. minae*, *P. pyrgillospora* and *P. xanthominuta* which are keyed out via 142 because they have a thick outer wall suggesting an apical endospore layer. Their ascospores resemble *Pyrgillus javanicus*)
- B121a Thallus UV+yellow. 122
- B121b Thallus UV-negative. 124
- B122a Hamathecium not inspersed; pantropical. *Pyrenula cocoes* Müll. Arg.
- B122b Hamathecium inspersed. 123
- B123a Thallus olivaceous with thick cortex; northern temperate. *Pyrenula pseudobufonia* (Rehm) R.C. Harris
- B123b Thallus pale grey, with thin cortex; neotropical (Guyana, Venezuela). *Pyrenula guyanensis* Sipman & Aptroot
(*Pyrenula obvoluta* (Nyl.) R.C. Harris may be an older name, but in the lectotype the fertile warts are only 0.6–0.8 mm wide and details of hamathecium and ascospores are missing)
- B124a Hamathecium inspersed (in some cases only in the upper part). 125
- B124b Hamathecium not inspersed. 136
- B125a Ascospores with elongated end locules. 126
- B125b Ascospores with all locules more or less rounded to angular. 129

- B126a Ascospores 20–25 μm long; pantropical (compare also *Pyrenula acutalis* which is only inspersed in the upper part). *Pyrenula maravalensis* Vain.
B126b Ascospores >25 μm long. 127
- B127a Ascospores with attenuated, horn-like apices, (21–)25–29(–32) μm long; Brazil. *Pyrenula cornutispora* Aptroot & M. Cáceres
B127b Ascospores without horn-like apices. 128
- B128a Ascospores 26–30 μm long; Australasian (Papua New Guinea). *Pyrenula rinodinospora* Aptroot
B128b Ascospores 35–50 μm long; eastern palaeotropical (India). *Pyrenula kurzii* Ajay Singh & Upreti
- B129a Ascospores all <16 μm long; pantropical. *Pyrenula cayennensis* Müll. Arg.
B129b Ascospores partly >16 μm long. 130
- B130a Ascospores <40 μm long. 132
B130b Ascospores >40 μm long. 135
- B131a Ascospores partly >30 μm long. 132
B131b Ascospores all <30 μm long. 133
- B132a Ascospores with black granular material around the ends; Australasian (Papua New Guinea). *Pyrenula grossa* Aptroot
B132b Ascospores without black granular material; eastern palaeotropical (Borneo, Sri Lanka). *Pyrenula fulva* (Krempelh.) Müll. Arg.
- B133a Thallus with patches of orange brown colouration which are KOH+purple; known from temperate regions on the northern hemisphere and South Africa, so probably cosmopolitan. *Pyrenula occidentalis* (R.C. Harris) R.C. Harris
B133b Thallus without coloured, KOH+purple patches. 134
- B134a Hamathecium only inspersed in the upper part; neotropical. *Pyrenula acutalis* R.C. Harris
B134b Hamathecium totally inspersed; pantropical. *Pyrenula fetivica* (Krempelh.) Müll. Arg.
- B135a Ascospores 42–50 μm long, mostly straight; eastern palaeotropical (India). *Pyrenula subcamptospora* Upreti
B135b Ascospores 45–52 μm long, mostly curved; neotropical. *Pyrenula cryptostoma* (Nyl.) Müll. Arg.

- B136a End locules elongated. 137
B136b All locules more-or-less rounded to angular. 139
- B137a Ascospores 20–29 μm long; ascomata <0.7 mm diam. 138
B137b Ascospores 27–40 μm long; ascomata >0.7 mm diam.; pantropical. *Pyrenula vernicosa* (Krempelh.) Müll. Arg.
- B138a Ascospores 20–29 μm long; ascomata 0.5–0.7 mm diam.; eastern palaeotropical. *Pyrenula finitima* Müll. Arg.
B138b Ascospores 20–29 μm long; ascomata <0.5 mm diam.; pantropical. *Pyrenula approximans* (Krempelh.) Müll. Arg.
- B139a Ascomata >2 mm diam.; eastern palaeotropical (Philippines). *Pyrenula irosina* Vain.
B139b Ascomata <2 mm diam. 140
- B140a Ascospores 11–13 μm long; palaeotropical (Seychelles). *Pyrenula infracongruens* Aptroot & Schumm
B140b Ascospores >13 μm long. 141
- B141a Ascospores 14–16 μm long; eastern palaeotropical. *Pyrenula mastophoriza* (Nyl.) Zahlbr.
B141b Ascospores at least partly >16 μm long; pantropical. *Pyrenula nitidula* (Bres.) R.C. Harris
- B142a Hamathecium inspersion. 143
B142b Hamathecium not inspersion. 162
- B143a Hamathecium inspersion with coloured particles. 144
B143b Hamathecium inspersion with colourless oil globules. 146
- B144a Hamathecium inspersion with orange, K-negative crystals; Sri Lanka. *Pyrenula multicolorata* Weerakoon & Aptroot
B144b Hamathecium inspersion with red or yellow oil globules. 145
- B145a Hamathecium inspersion red; ascospores 13–16 μm long; Venezuela. *Pyrenula rubroinspersa* Aptroot & Sipman
B145b Hamathecium inspersion yellow; ascospores 29–35 μm long; Ecuador. *Pyrenula flavoinspersa* Aptroot & Sipman

- B146a Central ascospore locules elongated; Northern temperate. *Pyrenula subelliptica* (Tuck.) R.C. Harris
- B146b Central ascospore locules transversely lenticular to rounded. 147
- B147a Ascospores constricted at the central septum; little endospore thickening between the outer lumina and the wall; Costa Rica. *Pyrenula minae* Aptroot & Lücking
- B147b Ascospores not constricted, at most becoming constricted when postmature. 148
- B148a Ascomata mostly <0.7 mm diam. 149
- B148b Ascomata mostly >0.7 mm diam. 153
- B149a Ascospores 13–18 µm long; neotropical. *Pyrenula laetior* Müll. Arg.
- B149b Ascospores >18 µm long. 150
- B150a Ascospores about 18–25 µm long. 151
- B150b Ascospores >27 µm long. 152
- B151a Ascospores 18–20 µm long; thallus and ascomata without any anthraquinone; eastern palaeotropical (Singapore). *Pyrenula subglabrata* (Nyl.) Müll. Arg.
- B151b Ascospores 18–25 µm long; thallus and/or ascomata with patches of rusty anthraquinone; known from temperate regions on the northern hemisphere and South Africa, so probably cosmopolitan. *Pyrenula occidentalis* (R.C. Harris) R.C. Harris
- B152a Ascospores 25–31 µm long; thallus with pseudocyphellae; Brazil. *Pyrenula supralaetior* C.O. Mendonça, Aptroot & M. Cáceres
- B152b Ascospores 27–50 µm long; thallus without pseudocyphellae; eastern palaeotropical (India, Sri Lanka). *Pyrenula oculata* Ajay Singh & Upreti
- B153a Ascospores mostly 10–20 µm long. 154
- B153b Ascospores mostly >20 µm long. 156
- B154a Ascospores mostly 10–14 µm long, up to 4 µm wide; Brazil. *Pyrenula inframamillana* Aptroot & M. Cáceres
- B154b Ascospores mostly >15 µm long. 155
- B155a Ascospores mostly 14–17 µm long and about half as wide, biseriate in the ascus; pantropical. *Pyrenula castanea* (Eschw.) Müll. Arg.
- B155b Ascospores mostly 17–21 µm long and about one third as wide, uniseriate in the ascus; pantropical *Pyrenula mamillana* (Ach.) Trevis.
- B156a Ascospores <30 µm long. 157

- B156b Ascospores 30–45 µm long. 159
- B157a Ascoma chamber deeply immersed in the bark; only ostioles visible, often with red-brown pigment; Brazil. *Pyrenula celaticarpa* Aptroot & M. Cáceres
- B157b Ascomata erumpent, at least partially visible as black dots. 158
- B158a Ascospores rounded, uniseriate in the ascus; pantropical. *Pyrenula massariospora* (Starb.) R.C. Harris
- B158b Ascospores at least at one end pointed, biseriate in the ascus; neotropical. *Pyrenula acutalis* R.C. Harris
- B159a Ascomata conical, emergent. 160
- B159b Ascomata globose to hemiglobose. 161
- B160a Ascomata conical, emergent, sides spreading; eastern palaeotropical. *Pyrenula interducta* (Nyl.) Zahlbr.
- B160a Ascospores without black granular material, and with thin apical walls bulging at the spore tips; palaeotropical (Seychelles). *Pyrenula supracongruens* Aptroot & Schumm
- B161a Ascomata globose, immersed; neotropical. *Pyrenula mastophoroides* (Nyl.) Zahlbr.
- B161b Ascomata hemispherical, prominent; eastern palaeotropical. *Pyrenula ocellulata* Wijeyaratne, Lücking & Lumbsch
- B162a Thallus UV+ yellow (when material well preserved, sometimes only part of the thallus reacting). 163
- B162b Thallus UV-negative or greenish/whitish reflecting. 167
- B163a Ascospores mostly >24 µm long. 164
- B163b Ascospores mostly <24 µm long. 165
- B164a Ascospores 25–40 µm long, without black granules; Costa Rica (also reported from China). *Pyrenula andina* Aptroot
- B164b Ascospores 36–45 µm long, with black granules at the tips; temperate America. *Pyrenula caryae* R.C. Harris
- B165a Thallus brownish with thick cortex; ascomata not in distinct thalus warts. 166
- B165b Thallus whitish with thin cortex; ascomata immersed in distinct thallus warts; ascospores 15–22 µm long, ellipsoidal, not pointed; neotropical. *Pyrenula micromma* (Mont.) Trevis.

- (*Pyrenula obvoluta* (Nyl.) R.C. Harris & Aptroot is most probably a separate species, characterized by not-pointed ascospores)
- B166a Ascospores 9–11 μm long, ellipsoid; pantropical (Australia, Guyana). *Pyrenula xanthominuta* Aptroot
- B166b Ascospores 14–21 μm long, fusiform, pointed; cosmopolitan. *Pyrenula dermatodes* (Borrer) Schaer.
(somewhat variable, especially in UV-reaction, but always with pointed ascospores and thallus at least partly covering the ascomata. *Pyrenula shirabeicola* Kurok. & S. Nakan. is probably a further synonym. The oldest name found for specimens that are not reacting with UV is *Pyrenula subtrahens* (Nyl.) Müll. Arg.)
- B167a Ascospores mostly $>25 \mu\text{m}$ long. 168
- B167b Ascospores mostly $<25 \mu\text{m}$ long. 181
- B168a Ascospores 36–45 μm long, with or without black granules at the tips. 169
- B168b Ascospores $<40 \mu\text{m}$ long, without black granules at the tips. 170
- B169a Ascospores 36–45 μm long, with black granules at the tips; temperate America. *Pyrenula caryae* R.C. Harris
- B169b Ascospores 36–45 μm long, without black granules at the tips; pantropical. *Pyrenula subducta* (Nyl.) Müll. Arg.
- B170a End locules elongated, ascospores 24–29 μm long; Virgin Islands. *Pyrenula expectata* R.C. Harris
- B170b End locules not elongated. 171
- B171a Growing on rock, ascospores with dark bands between the locules, 25–30 μm long; Brazil. *Pyrenula quarzitica* Aptroot
- B171b Growing on bark, ascospores without dark bands. 172
- B172a Ascomata mostly $>0.7 \text{ mm}$ diam. 173
- B172b Ascomata mostly $<0.7 \text{ mm}$ diam. 176
(*Pyrenula lilacina* with lilac-colored thallus keys out here when the pigment is not observed)
- B173a Thallus with pseudocyphellae; temperate regions of the old world. *Pyrenula macrospora* (Degel.) Coppins & P. James
- B173b Thallus without pseudocyphellae. 174

- B174a Ascospores 27–32 µm long, when mature dark brown and constricted at the central septum instead of pale grey brown and constricted at the locules; Guyana. *Pyrenula minoides* Aptroot & Sipman
- B174b Ascospores pale greybrown and constricted at the locules. 175
- B175a Ascospores 24–29 µm long, thallus thick; eastern palaeotropical (Papua New Guinea, Juan Fernandez). *Pyrenula media* Aptroot
- B175b Ascospores 29–40 µm long, thallus thin; eastern palaeotropical. *Pyrenula complanata* (Mont.) Trevis.
- B176a Ascospores with diamond-shaped locules; temperate regions on both hemispheres. *Pyrenula chlorospila* (Nyl.) Arnold
- B176b Ascospores with somewhat rounded or quadrangular locules. 177
- B177a Ascospores with dark granules between the locules; temperate America and Japan. *Pyrenula macounii* R.C. Harris
- B177b Ascospores without dark bands. 178
- B178a Ascospores 32–42 µm long; temperate America and Japan. *Pyrenula punctella* (Nyl.) Trevis.
- B178b Ascospores mostly 25–37 µm long; tropical. 179
- B179a Central locules of ascospores much longer than wide; ascomata superficial, hemispherical; Guyana. *Pyrenula perfecta* Aptroot & Sipman
- B179b Central locules of ascospores not longer than wide. 180
- B180a Central locules much wider than long, ascomata conical, emergent, thallus without pseudocyphellae; pantropical. *Pyrenula mastophora* (Nyl.) Müll. Arg.
- B180b Central locules more-or-less rounded; ascomata somewhat rounded, often partly immersed in the thallus; thallus often with (generally sparse) pseudocyphellae; pantropical. *Pyrenula quassiicola* (Fée) Fée
- B181a Ascospores mostly 21–25 µm long. 182
- B181b Ascospores mostly <21 µm long. 189
- B182a Ascomata c. 3–4 mm diam.; Japan. *Pyrenula gigas* Zahlbr.
- B182b Ascomata <3 mm diam. 183
- B183a Ascomata with red, KOH+purple crystals inside. 184
- B183b Ascomata without red crystals. 185

-
- B184a Ascomata <0.5 mm diam.; temperate regions of Europe and Asia. *Pyrenula nitidella* (Flörke ex Schaer.) Müll. Arg.
- B184b Ascomata >0.5 mm diam.; temperate regions of Europe and Asia. *Pyrenula nitida* (Weigel) Ach.
- B185a Ascospores with angular diamond-shaped locules; temperate North and South America. *Pyrenula micheneri* R.C. Harris
- B185b Ascospores with rounded or quadrangular locules. 186
- B186a Ascospores with bands of dark granules between the locules, without much endospore thickening between the outer lumina and the wall; pantropical. *Pyrenula pyrgillospora* Aptroot
- B186b Ascospores without bands of dark granules. 187
- B187a Thallus with dense pseudocyphellae; postmature ascospores longitudinally plicate; Chile. *Pyrenula maritima* Sipman & Aptroot
- B187b Thallus without pseudocyphellae; ascospores not regularly plicate when postmature. 188
- B188a Ascospores with at least one pointed end; temperate northern hemisphere, extending to the tropics. *Pyrenula acutispora* Kalb & Hafellner
- B188b Ascospores with rounded ends; eastern palaeotropical (India). *Pyrenula submastophora* Ajay Singh & Upreti
- B189a Ascospores 7–10 µm long; Iran. *Pyrenula minutissima* Aptroot, Valadbeigi & Sipman
- B189b Ascospores >10 µm long. 190
- B190a Ascospores mostly <15 µm long. 191
- B190b Ascospores mostly >15 µm long. 195
- B191a Ascomata >0.7 mm diam.; pantropical. *Pyrenula atropurpurea* (Eschw.) Müll. Arg.
- B191b Ascomata <0.7 mm diam. 192
- B192a Ascospores 6–8 µm wide; pantropical. *Pyrenula brunnea* Fée
- B192b Ascospores 3–6 µm wide. 193
- B193a Ascospores 3.5–4.0 µm wide, rhomboidal; Brazil. *Pyrenula rhomboidea* Aptroot & M. Cáceres
- B193b Ascospores 4–6 µm wide. 194

- B194a Ascospores 4–6 µm wide; pantropical. *Pyrenula aspistea* (Ach.) Ach.
B194b Ascospores 5–6 µm wide; like *P. aspistea* but with immersed ascomata; Brazil.
Pyrenula minutispora Aptroot & M. Cáceres
- B195a Thallus at least partly 0.3 mm thick. 196
B195b Thallus thinner. 197
- B196a Ascospores at least partly constricted at the septa; Australasian (Papua New Guinea). *Pyrenula montana* Aptroot
B196b Ascospores not constricted; neotropical (Puerto Rico, Brazil). *Pyrenula psoriformis* Zahlbr.
- B197a Ascomata >0.7 mm diam. 198
B197b Ascomata <0.7 mm diam. 201
- B198a Thallus with pseudocyphellae; New Zealand. *Pyrenula deliquescens* (C. Knight) Müll. Arg.
B198b Thallus without pseudocyphellae. 199
- B199a Ascospores pale reddish brown instead of pale grey-brown; Brazil. *Pyrenula mattickiana* Sipman & Aptroot
B199b Ascospores pale grey-brown. 200
- B200a Locules rounded; eastern palaeotropical (India). *Pyrenula scutata* (Stirt.) Zahlbr.
B200b Locules angular; pantropical. *Pyrenula balia* (Krempelh.) R.C. Harris
- B201a Ascospores with distinct inner endospore layer around locules; neotropical (Florida, Louisiana). *Pyrenula rubrostoma* R.C. Harris
B201b Ascospores without extra endospore layers. 202
- B202a Ascospores with dark bands between the locules; North America and Asia.
Pyrenula confoederata R.C. Harris
B202b Ascospores without dark bands. 203
- B203a Ascomata <0.4 mm diam.. 204
B203b Ascomata reaching >0.4 mm diam.; pantropical (UV-negative material of *Pyrenula dermatodes* keys out here as well; The oldest name found for such specimens that are not reacting with UV is *Pyrenula subtrahens* (Nyl.) Müll. Arg.). *Pyrenula aggregata* (Fée) Fée
- B204a Thallus pale grey, with thin cortex and filled with crystals, not oily or olivaceous; Brazil. *Pyrenula albonigra* Aptroot, D.S. Andrade & M. Cáceres

B204b Thallus olivaceous, oily. 205

B205a Ascospores 18–20 µm long; palaeotropical. *Pyrenula tristissima* Vain.

B205b Ascospores <18 µm long; neotropical. *Pyrenula minor* Fée

Key C. Pyrenocarpous lichens with simple paraphyses and septate, hyaline, distoseptate ascospores

(*Celothelium*, *Lithothelium*; temperate *Celothelium* species are not keyed out as they are dubious; all saxicolous *Lithothelium* species are keyed out here as well to make the treatment of all Pyrenulaceae complete, even if some of them have brown ascospores)

C1a Ascospores filiform. *Celothelium*. 2

C1b Ascospores ellipsoid to fusiform. *Lithothelium*. 7

C2a Ascomata sideways fused. 3

C2b Ascomata solitary or with joined ostioles. 4

C3a Ascomata sideways grouped in dense, often linear, clusters; neotropical. *Celothelium cinchonarum* (Vain.) Vain.

C3b Ascomata in small groups; temperate. *Celothelium ischnobelum* (Nyl.) M.B. Aguirre

C4a Ostiole lateral to eccentric, fused or not. 5

C4b Ostiole apical, ascomata solitary. 6

C5a Ascomata less than 0.5 mm wide; neotropical. *Celothelium aciculiferum* (Nyl.) Vain.

C5b Ascomata over 0.5 mm wide; Korea. *Celothelium longisporum* K.H. Moon & Aptroot

(here also keys out from India: *Lithothelium filisporum* (Patw., Makhija & D. Rane) Kr.P. Singh & G.P. Sinha, which could well be a *Celothelium*)

C6a Without yellow pigment; neotropical. *Celothelium dominicanum* (Müll. Arg.) Aguirre

C6b With yellow pigmentation; Bermuda. *Celothelium lutescens* F. Berger & Aptroot

C7a Ascospores colourless. 8

C7b Ascospores red-brown to brown; on rock (species on bark are in keys A and B). 25

C8a Thallus on bark; ascospores 3-septate. 9

C8b Thallus on rock. 21

- C9a Ascospores 4/ascus; Thailand. *Lithothelium quadrisporum* Aptroot
C9b Ascospores mostly 8/ascus. 10
- C10a Ascomata with fused ostioles only or with fused ostioles and fused walls. 11
C10b Ascomata solitary. 14
- C11a Ascomata with fused ostioles only; thallus epiphloeodic, brownish to green; ascospores 12–15(–18) × 4–6 µm; pantropical. *Lithothelium obtectum* (Müll. Arg.) Aptroot
C11b Ascomata with fused ostioles and fused walls. 12
- C12a Thallus endophloeodic, brownish to grey; ascospores 18–23 × (6–)7–10 µm; pantropical. *Lithothelium illotum* (Vain.) Aptroot
C12b Thallus epiphloeodic, brownish to green. 13
- C13a Ascospores 18–23 × (6–)8–11 µm; neotropical. *Lithothelium paraguayense* Müll. Arg.
C13b Ascospores 16–18 × 7–8 µm; neotropical. *Lithothelium fugax* (Müll. Arg.) Aptroot
- C14a Thallus epiphloeodic, brownish green to greenish. 15
C14b Thallus endophloeodic, whitish to brownish grey. 17
- C15a Ascospores 10–12 × 4–4.5 µm; Florida. *Lithothelium microsporium* R.C. Harris
C15b Ascospores larger. 16
- C16a Ascospores 16–19 × 5.5–6.5 µm; South Africa. *Lithothelium almbornii* Aptroot
C16b Ascospores 27–37 × 12–17 µm; pantropical (Brazil, China). *Lithothelium grossum* Aptroot
- C17a Hymenium interspersed with oil droplets; ascospores (9–)10–14(–17) × (3–)4–5 µm; Japan. *Lithothelium japonicum* H. Harada
C17b Hymenium not interspersed. 18
- C18a Terminal lumina much smaller than the central lumina. 19
C18b Terminal lumina only slightly smaller than the central lumina. 20
- C19a Ascospores 14–20(–24) × 6–8(–12) µm; pantropical. *Lithothelium hyalosporum* (Nyl.) Aptroot
C19b Ascospores 18–31 × 10–20 µm; Australia. *Lithothelium quiescens* P.M. McCarthy
- C20a Paraphyses simple; ascospores 18–23 × (6–)7–10 µm; pantropical. *Lithothelium illotum* (Vain.) Aptroot

- C20b Paraphyses branched and anastomosing; ascospores $17\text{--}25 \times 8.5\text{--}12 \mu\text{m}$; India. *Lithothelium insulare* Makhija & Adaw.
- C21a Ascospores 4–8-septate, $20\text{--}25(-30) \times 3\text{--}5.5(-6) \mu\text{m}$; Bahamas. *Lithothelium bahamense* Riddle
- C21b Ascospores 3-septate. 22
- C22a Ascomata with fused ostioles and fused walls; thallus endolithic. 23
- C22b Ascomata solitary; thallus at least partly epilithic. 24
- C23a Thallus violet-grey; hamathecium IKI+ blue to orange; asci with rounded ocular chamber; ascospores $13\text{--}17(-20) \times 3.5\text{--}5(-6) \mu\text{m}$; neotropical. *Lithothelium cubanum* Müll. Arg.
- C23b Thallus greenish grey; hamathecium IKI negative; asci with sagittiform ocular chamber; ascospores $16\text{--}19 \times 5\text{--}7 \mu\text{m}$; Falklands. *Lithothelium falklandicum* (Nyl.) Aptroot
- C24a Ostioles apical; asci without ocular chamber; ascospores $17\text{--}20 \times 6\text{--}8 \mu\text{m}$; Eurasian temperate. *Lithothelium triseptatum* (Nyl.) Aptroot
- C24b Ostioles lateral; asci with rounded ocular chamber; ascospores $15\text{--}18 \times 6\text{--}7 \mu\text{m}$; Hawaii. *Lithothelium hawaiiense* Aptroot
- C25a Ascospores ornamented with spines, $14\text{--}19 \times 6\text{--}7.5 \mu\text{m}$; China. *Lithothelium echinatum* Aptroot
- C25b Ascospores smooth. 26
- C26a Thallus epilithic, rimose to areolate; ascospores $18\text{--}28 \times 8\text{--}14 \mu\text{m}$; Australia. *Lithothelium austropacificum* P.M. McCarthy
- C26b Thallus endolithic; ascospores $20\text{--}26 \times 6\text{--}8 \mu\text{m}$. 27
- C27a Ascospore lumina all rounded: New Zealand. *Lithothelium australe* Aptroot & H. Mayrhofer
- C27b Ascospore lumina partly diamond-shaped and connected; Bermuda. *Lithothelium bermudense* F. Berger, LaGreca & Aptroot

Key D. Tropical pyrenocarpous lichens with simple paraphyses and 1-septate, distoseptate, brown ascospores (mostly Pyrenulaceae: *Clypeopyrenis*, *Distopyrenis*, *Granulopyrenis*, *Parapyrenis*)

(all are rare or rarely collected and rarely studied; at least the sequenced *Granulopyrenis* is basal in or sister to the Pyrenulaceae)

D1a ascomata conical, distinctly flattened, superficial, 0.6–1.2 mm diam.; ostioles apical, black, obconical; ascospores brown, biloculate, $6-8 \times 2-2.5 \mu\text{m}$, ellipsoidal; neotropical. *Clypeopyrenis microsperma* (Müll. Arg.) Aptroot

D1b Ascospores larger. 2

D2a Ascospores with double wall, with a layer of conspicuous granules inside.

Granulopyrenis. 3

D2b Ascospores without internal granules. 7

D3a Ascomata grouped under a stroma; Brazil. *Granulopyrenis nigrescens* (Müll. Arg.) R.C. Harris

D3b Ascomata single. 4

D4a Thallus not blackened around the ascocarps; ostiole apical, asci with rounded ocular chamber; South Africa. *Granulopyrenis macrocarpoides* (Zahlbr.) Aptroot

D4b Thallus blackened around the ascocarps. 5

D5a Ostiole lateral; asci with rounded ocular chamber; on bark; on driftwood; USA.

Granulopyrenis hymnothora Aptroot

D5b Ostiole apical to eccentric; asci without ocular chamber. 6

D6a Ostiole partly eccentric; Caribbean. *Granulopyrenis portoricensis* (Petrač) Y. Z. Wang, Aptroot & K. D. Hyde

D6b Ostiole apical; Hong Kong. *Granulopyrenis seawardii* Aptroot & Sipman

D7a Thallus lichenicolous; Papua New Guinea. *Parapyrenis lichenicola* Aptroot & Diederich

D7b Thallus not lichenicolous. 8

D8a Thallus on driftwood; Papua New Guinea. *Parapyrenis maritima* Aptroot

D8b Thallus on bark. 9

D9a Ascomata grouped under a stroma; neotropical (Brazil, Puerto Rico). *Distopyrenis composita* R.C. Harris

D9b Ascomata single. 10

D10a Thallus with violet tinge; Australia. *Distopyrenis violacea* Aptroot

D10b Thallus brownish or whitish. 11

- D11a Ascospores with elongated lumina. 12
 D11b Ascospores with rounded or angular lumina. 13
- D12a hamathecium IKI + blue; ascocarps flattened, simple or aggregated; Bermuda.
Parapyrenis elongata Aptroot
 D12b hamathecium IKI negative; ascocarps not flattened, simple; neotropical.
Parapyrenis guayaca (Fée) Aptroot
- D13a Ascospores with an euseptum within the septum. 14
 D13b Ascospores only distoseptate with relatively small lumina. 15
- D14a Lumina angular; thallus whitish grey to invisible, not blackened around the
 ascocarps; USA and Caribbean. *Distopyrenis americana* Aptroot
 D14b Lumina rather rounded; thallus pinkish grey, blackened around the ascocarps;
 Mexico and Caribbean. *Distopyrenis fuliginosa* Aptroot
- D15a Ascospores with large, rounded lumina; Australia. *Parapyrenis conica* Aptroot
 D15b Ascospores with angular lumina, with much thickened endosporium formation,
 thus lumina relatively small. 16
- D16a Ascomata flattened; Java. *Parapyrenis aurora* (Zahlbr.) Aptroot
 D16b Ascomata conical; French Guiana. *Distopyrenis pachyspora* Aptroot

**Key E. Pyrenocarpous lichens with mazaedioid ascomata (Pyrenulaceae:
Mazaediothecium, Pyrgillus)**

- E1a Ascoma wall not carbonized; ascospores amassing in the large ostioles of the cup-
 like ascomata, either 1-septate or cruciate submuriform. *Mazaediothecium*. 2
 E1b Ascoma wall carbonized; ascospores amassing in a cup-like structure on top of the
 closed ascomata, transversely 1-3-septate. *Pyrgillus*. 6
- E2a Ascomata red; Papua New Guinea. *Mazaediothecium rubiginosum* Aptroot
 E2b Ascomata black. 3
- E3a Ascospores 1-septate. 4
 E3b Ascospores cruciate submuriform. 5
- E4a Thallus corticate; photobiont trebouxoid; Brazil. *Mazaediothecium serendipiticum*
 Aptroot

-
- E4b Thallus not corticate; photobiont trentepohlioid; French Guiana. *Mazaediothecium uniseptatum* Aptroot
- E5a Thallus UV+ yellow; neotropical. *Mazaediothecium album* Aptroot
 E5b Thallus UV-negative; Malaysia. *Mazaediothecium mohamedii* H. Harada & Yoshik. Yamam.
- E6a Ascospores 1-septate. 7
 E6b Ascospores 3-septate. 9
- E7a Mazaedium pruina golden-yellow; pantropical. *Pyrgillus indicus* (Kremp.) Aptroot
 E7b Mazaedium pruina white. 8
- E8a Ascospores with pale ends, red brown; Australia. *Pyrgillus fuscus* Aptroot
 E8b Ascospores with dark ends, brown; pantropical. *Pyrgillus cubanus* Nyl.
- E9a Thallus UV–; mazaedium pruina orange, KOH+ purple; ascospores 10–14 × 5–7 µm; Cambodia and China. *Pyrgillus cambodiensis* Kashiw., K.H. Moon & Aptroot
 E9b Thallus UV+ yellow, with lichexanthone. 10
- E10a Mazaedium pruina white, KOH–; ascospores 10–13 × 6.0–8.5 µm; pantropical. *Pyrgillus javanicus* (Mont. & Bosch) Nyl.
 E10b Mazaedium pruina orange or red, KOH+ orange, reddish violet or violet. 11
- E11a Mazaedium pruina red, KOH+ reddish violet, UV–; ascospores 15–20(–23) × 7.5–10.0 µm; India. *Pyrgillus tibellii* Kr. P. Singh & Pushpi Singh
 E11b Mazaedium pruina red but KOH+ violet or orange, UV+ red; ascospores 13.0–17.5 × 5.0–7.5 µm. 12
- E12a Mazaedium pruina orange, KOH+ violet; ascospores 13–16 × 6.0–7.5 µm, c. 2 times as long as wide; Brazil. *Pyrgillus aurantiacus* Aptroot & M. Cáceres
 E12b Mazaedium pruina red, KOH+ orange; ascospores 15.0–17.5 × 5.0–6.5 µm, c. 2.5–3 times as long as wide; Brazil. *Pyrgillus rufus* Aptroot & M. Cáceres

Key F. *Aptrootia* species

- F1a Thallus on bark, corticate, green, bullate; ascospores ornamented; Eastern palaeotropical. *Aptrootia elatior* (Stirt.) Aptroot

F1b Thallus on soil and bryophytes, not corticate, grey, cartilaginous, smooth; ascospores smooth. 2

F2a Ascospores $170\text{--}230 \times 40\text{--}70 \mu\text{m}$; amphipacific (Costa Rica, Papua New Guinea, Solomon Islands). *Aptrootia terricola* (Aptroot) Lücking, L. Umaña & Chaves

F2b Ascospores $150\text{--}360\text{--}400 \times 65\text{--}140 \mu\text{m}$; Australia. *Aptrootia robusta* (P.M. McCarthy & Kantvilas) Aptroot

Key G. *Architrypethelium* species with transversely septate ascospores

G1a Ascospores remaining hyaline 2

G1b Ascospores soon becoming brown 5

G2a Ascomata sessile, very large (up to 3 mm diam.); Brazil. *Architrypethelium grande* (Kremp.) Aptroot & Lücking

G2b Ascomata immersed or in erumpent prominent thallus warts, smaller (usually not exceeding 1 mm diam.). 3

G3a Ascomata completely immersed between bullate thallus warts; Brazil. *Architrypethelium lauropaluanum* Lücking, M.P. Nelsen & Marcelli

G3b Ascomata erumpent to prominent, partly thallus-covered. 4

G4a Thallus UV+ yellow; neotropical. *Architrypethelium hyalinum* Aptroot

G4b Thallus UV–; Bolivia. *Architrypethelium penuriixanthum* Flakus & Aptroot

G5a Ostiole eccentric; ascomata sometimes with fused ostioles; neotropical. *Architrypethelium nitens* (Fée) Aptroot

G5b Ostiole apical; ascomata solitary. 6

G6a Hamathecium inspersed; Colombia. *Architrypethelium columbianum* (Nyl.) Aptroot & Lücking

G6b Hamathecium clear; pantropical. *Architrypethelium uberinum* (Fée) Aptroot

Key H. *Astrothelium* species with ascospores transversely septate, ostiole apical, secondary substances present on thallus and/or ascoma

H1a Lichexanthone present; thallus and/or ascomata UV+ yellow; pigment(s) present or absent. 2

- H1b Lichexanthone absent; pigment(s) always present, thallus and ascomata usually K+ red to purple and UV+ orange or red. 20
- H2a Pigments present in thallus and/or ascomata; lichexanthone on thallus; ascomata pseudostromatic, in irregular black lines. 3
- H2b Pigments absent; lichexanthone on thallus and/or ascomata; ascomata variable. 4
- H3a Ascospores 3-septate, small (18–26 μm long); ascomata with external orange pigment, scattered laterally along pseudostromata; neotropical. *Astrothelium neogalbineum* (R.C. Harris) Aptroot & Lücking
- H3b Ascospores (3–)5-septate, large (90–115 μm long); ascomata with external purplish-red pigment, scattered laterally along pseudostromata; Brazil. *Astrothelium pictum* Aptroot
- H4c Ascospores 9–11-septate; Brazil. *Astrothelium pseudodermatodes* Aptroot
- H4a Ascospores 3–9-septate. 5
- H5a Ascospores 5–9-septate. 6
- H5b Ascospores 3-septate. 7
- H6a Lichexanthone on ascomata; ascomata solitary to irregularly confluent, erumpent, covered by thallus; eastern palaeotropical. *Astrothelium cinereorosellum* (Kremp.) Aptroot & Lücking
- H6b Lichexanthone on pseudostromata; ascomata distinctly pseudostromatic; Fiji. *Astrothelium fijiense* Lücking, Nakswankul & Lumbsch
- H7a Hamathecium inspersed; ascospores small (15–30 μm long). 8
- H7b Hamathecium clear; ascospores small to medium-sized (15–70 μm long). 12
- H8a Lichexanthone on ascomata only, absent from thallus; ascomata solitary to irregularly confluent, largely immersed and covered by thallus; Brazil. *Astrothelium punctulatum* Malme
- H8b Lichexanthone on thallus and ascomata or on thallus only. 9
- H9a Lichexanthone on thallus only, absent from ascomata; ascomata solitary but dense; thallus gall-like; eastern palaeotropical. *Astrothelium grossoides* Aptroot & Lücking
- H9b Lichexanthone on thallus and ascomata; thallus not gall-like. 10
- H10a Ascomata pseudostromatic, immersed in prominent pseudostromata that are often partly covered by thallus; pantropical. *Astrothelium porosum* (Ach.) Aptroot & Lücking

(the recently described *Astrothelium sinense* S.H. Jiang & C. Zhang keys out here and is probably synonymous)

H10b Ascomata solitary, erumpent, whitish. 11

H11a Ascospores 20–25 μm long; neotropical. *Astrothelium vulcanum* Aptroot

H11b Ascospores 14–17 μm long; Brazil. *Astrothelium infravulcanum* Aptroot

H12a Ascospores small (15–30 μm long). 13

H12b Ascospores (small to) medium-sized (25–70 μm long). 17

H13a Lichexanthone only on ascomata or ostiole. 14

H13b Lichexanthone on thallus and ascomata. 15

H14a Lichexanthone on ascomata only, absent from thallus; ascomata pseudostromatic, immersed in prominent pseudostromata covered by thallus; neotropical (Brazil and Bolivia). *Astrothelium stromatofluorescens* Aptroot & M. Cáceres

H14b Lichexanthone only on ostioles; pseudostromata sessile, brownish; Brazil. *Astrothelium gyalostiolum* Aptroot

H15a Ascomata solitary to irregularly confluent, erumpent, covered by thallus; pantropical. *Astrothelium pulcherrimum* (Fée) Aptroot & Lücking

H15b Ascomata pseudostromatic, immersed in pseudostromata that are often partly covered by thallus. 16

H16a Pseudostromata well-delimited, with whitish cover, prominent to sessile; neotropical. *Astrothelium leucosessile* Lücking, M.P. Nelsen & Aptroot

H16b Pseudostromata not well-delimited, covered by thallus, erumpent to prominent; pantropical. *Astrothelium phlyctaena* (Fée) Aptroot & Lücking

H17a Ascospores under 40 μm long. 18

H17b Ascospores over 45 μm long. 19

H18a Ascospores 25–40 \times 8–13 μm ; thallus and thalline cover of ascomata UV+ yellow; pantropical. *Astrothelium pupula* (Ach.) Aptroot & Lücking

H18b Ascospores 34–38 \times 8.5–10.5 μm ; only pseudocyphellae UV+ yellow; Brazil. *Astrothelium xanthopseudocyphellatum* Aptroot

H19a Ascospores 60–70 \times 16–18 μm ; El Salvador. *Astrothelium megochroleucum* Aptroot

H19b Ascospores 45–60 \times 19.5–22.5 μm ; Brazil. *Astrothelium miniannulare* Kalb & Aptroot

- H20a Ascospores 3-septate. 21
H20b Ascospores 5–19-septate. 44
- H21a Ascospores medium-sized (50–65 µm long); medulla (ascomata) with internal, redpigment; ascomata solitary to irregularly confluent, erumpent, covered by thallus; neotropical. *Astrothelium thelotremoides* (Nyl.) Aptroot & Lücking
H21b Ascospores small to medium-sized (15–50 µm long); pigment external or internal, yellow to red. 22
- H22a Hamathecium inspersed. 23
H22b Hamathecium clear. 28
- H23a Hamathecium with red, K+ green pigment; Sri Lanka. *Astrothelium isohypocrellinum* Aptroot & Weerakoon
H23b Hamathecium hyaline. 24
- H24a Ascomata with external, yellow to orange pigment mostly developed around the ostiolar area; ascomata solitary to irregularly confluent, erumpent, covered by thallus except for black ostiolar area. 25
H24b Medulla (ascomata) with internal, red pigment; ascomata variable. 27
- H25a Ascomata with yellow pigment; ascomata diffusely pseudostromatic, pseudostromata forming irregular lines; El Salvador. *Astrothelium neoinspersum* Aptroot
H25b Thallus and/or ascomata with orange pigment; ascomata solitary to irregularly confluent. 26
- H26a Both thallus and ascomata with abundant orange pigment; pantropical. *Astrothelium inspersaeneum* Aptroot & M. Cáceres
H26b Only ascomata with sparse orange pigment; pantropical (Colombia and Papua New Guinea). *Astrothelium aenascens* Aptroot
- H27a Ascospores 20–30 × 7–9 µm; Brazil. *Astrothelium rubrocrystallinum* Aptroot & M. Cáceres
H27b Ascospores 30–50 × 14–16 µm; neotropical. *Astrothelium buckii* (R.C. Harris) Aptroot & Lücking
- H28a Thallus and/or ascomata with external pigment(s). 29
H28b Medulla (ascomata) with internal pigment(s). 39
- H29a Ascomata with external, red pigment(s); Mexico. *Astrothelium coccineum* Córd.-Cháv., Aptroot & M. Cáceres

- H29b Ascomata with external, yellow to orange pigment(s). 30
- H30a Thallus grey, barely lichenized; ascomata distinctly pseudostromatic, pseudostromata prominent to sessile, with orange cover strongly contrasting with thallus; Thailand. *Astrothelium aurantiacocinereum* Lücking, Naksuwankul & Lumbsch
- H30b Thallus olive-green, distinctly lichenized; ascomata solitary or pseudostromatic, if pseudostromatic, then pseudostromata immersed to erumpent and in irregular, often reticulate lines. 31
- H31a Ascospores very small (16–18 µm long); ascomata pseudostromatic, immersed, pseudostromata in irregular, often reticulate lines, covered with pigment and contrasting with olive-green thallus; Brazil. *Astrothelium flavum* Aptroot & M. Cáceres
- H31b Ascospores small (20–35 µm long). 32
- H32a Ascospores 20–25(–27) × 6–10 µm; ascomata solitary or pseudostromatic. 33
- H32b Ascospores 25–35 × 10–12 µm; ascomata solitary to irregularly confluent, erumpent, covered by thallus. 37
- H33a Ascomata solitary to irregularly confluent; pigment on thallus and ascomata. 34
- H33b Ascomata pseudostromatic, pseudostromata in irregular lines, covered with pigment and contrasting with olive-green thallus. 36
- H34a Thallus resembling bubbles of wax plus prothallus; Venezuela. *Astrothelium bullatothallinum* Aptroot & Sipman
- H34b Thallus smooth. 35
- H35a Pigment orange to yellow; pantropical. *Astrothelium aeneum* (Eschw.) Aptroot & Lücking
- H35b Pigment ochraceous; Brazil. *Astrothelium ochraceum* Aptroot
- H36a Thallus strongly bullate; Brazil. *Astrothelium flavostromatum* Aptroot & M. Cáceres
- H36b Thallus smooth to uneven; neotropical. *Astrothelium kunzei* (Fée) Aptroot & Lücking
- H37a Pigment orange, abundant; neotropical (Brazil & Bolivia). *Astrothelium megaeneum* Flakus & Aptroot
- H37b Pigment yellow, sparse. 38

- H38a Ostioles partly lateral and fused; pigment K+ purple; Bolivia. *Astrothelium pallidoflavum* Flakus & Aptroot
- H38b Ostioles consistently apical; pigment K+ red; neotropical. *Astrothelium subcatervarium* (Malme) Aptroot & Lücking
- H39a Ascospores medium-sized (30–50 × 10–16 µm). 40
- H39b Ascospores small (15–30 × 5–9 µm). 42
- H40a Medullary pigment red; neotropical. *Astrothelium cartilagineum* (Fée) Aptroot & Lücking
- H40b Medullary pigment yellow. 41
- H41a Thallus bullate; Brazil. *Astrothelium endochryseum* (Vain.) Aptroot & Lücking
- H41b Thallus smooth; Brazil. *Astrothelium laevithallinum* Lücking, M.P. Nelsen & Marcelli
- H42a Ascomata solitary to irregularly confluent, erumpent, covered by thallus; medullary pigment orange; Brazil. *Astrothelium aeneoides* Aptroot
- H42b Ascomata pseudostromatic; pseudostromata blackish brown; medullary pigment (pale) yellow. 43
- H43a Medullary pigment K+ red; pantropical. *Astrothelium degenerans* (Vain.) Aptroot & Lücking
- H43b Medullary pigment K-. 16; neotropical. *Astrothelium feei* (C.F.W. Meissn.) Aptroot & Lücking
- H44a Ascospores small to medium-sized (15–45 × 6–9 µm). 45
- H44b Ascospores medium-sized to large (50–150 × 10–40 µm). 47
- H45a Medulla (ascomata) with internal, red pigment; ascospores 9–13-septate; ascomata pseudostromatic, exposed, forming irregular, reticulate lines or groups; India. *Astrothelium nigrorufum* (Makhija & Patw.) Aptroot & Lücking
- H45b Medulla (ascomata) with internal, yellow to orange pigment; ascospores 5–9-septate. 46
- H46a Hamathecium inspersed; medullary pigment K+ red; ascomata pseudostromatic, pseudostromata erumpent to prominent, whitish; Papua New Guinea. *Astrothelium pseudodissimulum* Aptroot
- H46b Hamathecium clear; medullary pigment K+ purple; ascomata pseudostromatic, exposed, forming irregular, reticulate lines or groups; India. *Astrothelium dissimilum* (Makhija & Patw.) Aptroot & Lücking

- H47a Ascospores medium-sized (50–90 μm long). 48
H44b Ascospores large (110–150 μm long). 49
- H48a Ascomata with scattered, external, pink pigment, solitary to irregularly confluent, immersed in large thallus verrucae; hamathecium inspersed; ascospores 50–65 \times 10–17 μm ; Brazil. *Astrothelium decemseptatum* Aptroot & M. Cáceres
H48b Medulla (ascomata) with internal, yellow pigment; ascomata pseudostromatic, pseudostromata prominent, covered by thallus except black ostiolar areas; hamathecium clear; ascospores 55–90 \times 13–28 μm ; East Asia *Astrothelium luridum* (Zahlbr.) Aptroot & Lücking
- H49a Ascomata with scattered, external, pink pigment, solitary, immersed; ascospores 17–22 μm broad, 13–19-septate; Caribbean. *Astrothelium calosporum* (Müll. Arg.) Aptroot & Lücking
H49b Ascomata with external and internal yellow pigment; ascospores 30–37 μm broad, 7–11-septate; neotropical. *Astrothelium rimosum* Aptroot

Key I. *Astrothelium* species with ascospores transversely septate, ostiole apical, secondary substances absent

- I1a Ascospores 3-septate. 2
I1b Ascospores (3–)5–15-septate. 26
- I2a Ascospores small to medium-sized (15–45(–50) \times 6–16 μm). 3
I2b Ascospores medium-sized to large ((45–)50–120 \times 15–35 μm). 20
- I3a Hamathecium inspersed. 4
I3b Hamathecium clear. 6
- I4a Ascospores (25–)30–40 μm long; ascomata in irregular groups to pseudostromatic, pseudostromata erumpent, white, contrasting with olive-green thallus; pantropical. *Astrothelium perspersum* Aptroot & Ertz
I4b Ascospores 15–25(–30) μm long. 5
- I5a Ascomata solitary, erumpent, with white cover; neotropical. *Astrothelium subscoria* Flakus & Aptroot
I5b Ascomata irregularly grouped to pseudostromatic, immersed to erumpent, mostly exposed, brown-black; pantropical. *Astrothelium scoria* (Fée) Aptroot & Lücking
- I6a Ascospores I+ violet. 7

- I6b Ascospores I–. 8
- I7a Ascomata solitary, immersed, covered by thallus except dark ostiolar area; pantropical. *Astrothelium infossum* (Nyl.) Aptroot & Lücking
- I7b Ascomata distinctly pseudostromatic, pseudostromata erumpent, brown-black, with pale ostiolar areas; neotropical. *Astrothelium infuscatulum* (Müll. Arg.) Aptroot & Lücking
- I8a Ascospores $15\text{--}28 \times 7\text{--}10\text{--}(12) \mu\text{m}$ (*Astrothelium nitidiusculum* complex). 9
- I8b Ascospores $26\text{--}46 \times (7\text{--})10\text{--}16 \mu\text{m}$. 15
- I9a Ascomata pseudostromatic; pseudostromata either prominent or contrasting with thallus in colour. 10
- I9b Ascomata solitary to irregularly confluent or grouped but not prominent or contrasting with thallus. 13
- I10a Pseudostromata prominent, partly covered by thallus and hence not strongly contrasting with thallus in colour; pantropical. *Astrothelium bicolor* (Taylor) Aptroot & Lücking
- I10b Pseudostromata immersed to erumpent and more or less leveled with thallus, often forming irregular to reticulate lines or groups, contrasting with thallus in colour (either dark or whitish). 11
- I11a Pseudostromata with whitish cover; neotropical. *Astrothelium rufescens* (Müll. Arg.) Aptroot & Lücking
- I11b Pseudostromata exposed, brown-black. 12
- I12a Pseudostromata forming distinct, compact groups laterally covered by thallus; neotropical. *Astrothelium subendochryseum* Lücking, M.P. Nelsen & Marcelli
- I12b Pseudostromata diffuse, exposed; neotropical. *Astrothelium nigratum* (Müll. Arg.) Aptroot & Lücking
- I13a Ascomata forming irregular groups, with rather broad, brown-black ostiolar area; eastern palaeotropical. *Astrothelium meiophorum* (Nyl.) Aptroot & Lücking
- I13b Ostiolar areas not broadly. 14
- I14a Ascomata covered by olive-green thallus; neotropical. *Astrothelium ceratinum* (Fée) Aptroot & Lücking
- I14b Ascomata with broadly white area surrounding the ostiole or entirely white; pantropical. *Astrothelium nitidiusculum* (Nyl.) Aptroot & Lücking

- I15a Ascospores $36\text{--}46 \times 14\text{--}16 \mu\text{m}$; ascomata solitary, erumpent, covered by thallus and with pale ring surrounding the dark ostiolar spot; pantropical, extending into temperate regions. *Astrothelium floridanum* Zahlbr. ex M. Choisy
- I15b Ascospores $26\text{--}39 \times 7\text{--}14 \mu\text{m}$; ascomata variable. 16
- I16a Ascomata pseudostromatic, pseudostromata exposed, immersed and leveled with thallus, forming irregular to reticulate lines or groups. 17
- I16b Ascomata solitary, covered by thallus. 19
- I17a Thallus consisting of globose verrucae with a thick hyaline cortex; Papua New Guinea. *Astrothelium diaphanocorticatum* Aptroot & Sipman
- I17b Thallus smooth to verrucose. 18
- I18a Ascospores $36\text{--}39 \mu\text{m}$ long; neotropical. *Astrothelium scoriiothelium* Aptroot & Lücking
- I18b Ascospores $29\text{--}33 \mu\text{m}$ long; Brazil. *Astrothelium disjunctum* Aptroot & M. Cáceres
- H19a Thallus and ascomata finely verrucose; ascomata immersed to erumpent, numerous and dense; neotropical. *Astrothelium papulosum* (Nyl.) Aptroot & Lücking
- H19b Thallus and ascomata smooth to uneven; ascomata erumpent to prominent, scattered; neotropical. *Astrothelium solitarium* Aptroot & M. Cáceres
- I20a Hamathecium inspersed. 21
- I20b Hamathecium clear. 23
- I21a Ascospores $45\text{--}62 \times 15\text{--}22 \mu\text{m}$; thallus uneven-bullate; ascomata solitary, erumpent, covered by thallus, with pale ring surrounding the dark ostiolar spot; Bolivia. *Astrothelium inspersotuberculosis* Flakus & Aptroot
- I21b Ascospores $85\text{--}120 \times 22\text{--}35 \mu\text{m}$; thallus and ascomata variable. 22
- I22a Ascospores $85\text{--}95 \times 22\text{--}25 \mu\text{m}$; thallus uneven-verrucose; ascomata solitary, prominent, exposed, with gently sloping sides; Vietnam. *Astrothelium clypeatum* Aptroot & Gueidan
- I22b Ascospores $100\text{--}120 \times 30\text{--}35 \mu\text{m}$; thallus uneven-verrucose; ascomata confluent, erumpent to prominent, exposed, with gently to steeply sloping sides; Guyana. *Astrothelium megatropicum* Aptroot
- I23a Ascomata prominent, completely exposed; Colombia. *Astrothelium phaeothelium* (Nyl.) Aptroot & Lücking
- I23b Ascomata immersed to erumpent, covered by thallus, with pale ring surrounding the dark ostiolar spot. 24

-
- I24a Thallus verrucose-papillose; ascomata solitary to irregularly confluent, prominent to sessile; Panama. *Astrothelium pseudannulare* Aptroot & Etayo
I24b Thallus uneven to shallowly bullate. 25
- I25a Ascospores 48–65 × 17–20 µm; neotropical. *Astrothelium tuberculosum* (Vain.) Aptroot & Lücking
I25b Ascospores 62–80 × 20–25 µm; neotropical. *Astrothelium annulare* (Fée) Aptroot & Lücking
- I26a Ascospores medium-sized (29–59(–80) × 8–17 µm). 27
I26b Ascospores large (85–220 × (8–)20–45 µm). 32
- I27a Hamathecium inspersed. 28
I27b Hamathecium clear. 29
- I28a Ascospores 50–59 µm long; thallus verrucose-bullate; Papua New Guinea. *Astrothelium galligenum* (Aptroot) Aptroot & Lücking
I28b Ascospores 40–49 µm long; thallus smooth to uneven; Sri Lanka. *Astrothelium nitidulum* Weerakoon & Aptroot
- I29a Ascomata at least mostly solitary. 30
I29b Ascomata in irregular groups to pseudostromatic, groups or pseudostromata erumpent to prominent. 31
- I30a Ascomata solitary to irregularly confluent, immersed to erumpent, with erumpent part exposed, blackish; Malaysia. *Astrothelium tryptelizans* (Nyl.) Aptroot & Lücking
I30b Ascomata solitary, immersed, not exposed; Brazil. *Astrothelium quintannulare* Aptroot
- I31a Pseudostromata whitish; eastern palaeotropical. *Astrothelium ubianense* (Vain.) Aptroot & Lücking
I31b Pseudostromata brownish; Brazil. *Astrothelium pseudoplatystomum* (Makhija & Patw.) Aptroot & Lücking
- I32a Hamathecium inspersed. 33
I32b Hamathecium. 38
- I33a Ascospores 9–15-septate, occasionally with a longitudinal septum in one or several cells. 34
I33b Ascospores 5–9-septate, lacking longitudinal septa. 35

- I34a Ascospores 120–195 µm long; India. *Astrothelium macrosporum* (Makhija & Patw.) Aptroot & Lücking
- I34b Ascospores 75–100 µm long; Malaysia. *Astrothelium taniaum* Aptroot & Sipman
- I35a Ascospores 180–220 × 40–45 µm, 5–9-septate; ascomata grouped in large thallus verrucae; Angola. *Astrothelium pustulatum* (Vain.) Aptroot & Lücking
- I35b Ascospores 100–170 × 29–40 µm, 5-septate; ascomata variable. 36
- I36a Ascospores curved; ascomata grouped; Brazil. *Astrothelium curvisporum* Aptroot & M. Cáceres
- I36b Ascospores straight; ascomata solitary to irregularly confluent. 37
- I37a Ascomata solitary, prominent to sessile, covered by thallus except dark ostiolar area; neotropical (Brazil and Guyana). *Astrothelium sipmanii* Aptroot
- I37b Ascomata solitary to irregularly confluent, erumpent, upper part exposed; Colombia. *Astrothelium pseudomegalophthalmum* Aptroot
- I38a Ascospores 140–190 µm long, 11–19-septate; ascomata solitary to irregularly confluent. 39
- I38b Ascospores 85–120 µm long, 3–17-septate; ascomata irregularly grouped to pseudostromatic. 40
- I39a Thallus verrucose-bullate; ascomata erumpent to prominent, covered by thallus except broad, blackened ostiolar area; Uruguay. *Astrothelium basilicum* (Kremp.) Aptroot & Lücking
- I39b Thallus smooth; ascomata immersed, completely covered by thallus; Argentina. *Astrothelium spectabile* (Aptroot & L.I. Ferraro) Aptroot & Lücking
- I40a Ascospores 20–33 µm broad, 3–7-septate. 41
- I40b Ascospores 8–16 µm broad, 9–17-septate; neotropical. *Astrothelium olivaceofuscum* (Zenker) Aptroot & Lücking
- I41a Thallus smooth to uneven; ascomata covered by thallus; Brazil. *Astrothelium trematum* Kalb & Aptroot
- I41b Thallus verrucose-bullate; ascomata exposed, brown-black. 42
- I42a Thallus verrucose-bullate; Brazil. *Astrothelium megalophthalmum* (Müll. Arg.) Aptroot & Lücking
- I42b Thallus squamulose; Brazil. *Astrothelium squamosum* Aptroot

Key J. *Astrothelium* species with ascospores transversely septate, ostiole lateral, secondary substances present on thallus and/or ascoma

- J1a Lichexanthone present; thallus and/or ascomata UV+ yellow; pigment(s) present or absent. 2
- J1b Lichexanthone absent; pigment(s) always present, thallus and ascomata usually K+ red to purple and UV+ orange or red. 38
- J2a Pigments present on/in thallus and/or ascomata; lichexanthone on thallus only. 3
- J2b Pigments absent; lichexanthone on thallus and/or ascomata. 10
- J3a Ascospores 9–11-septate, $50\text{--}75 \times 12\text{--}22 \mu\text{m}$; thallus with external, orange pigment (absent on ascomata); ascomata solitary to irregularly confluent, erumpent, covered by thallus except dark ostiolar area; Venezuela. *Astrothelium dicoloratum* Aptroot
- J3b Ascospores 3-septate, $21\text{--}55 \times 7\text{--}23 \mu\text{m}$; ascomata with external, yellow-orange pigment (absent on thallus). 4
- J4a Hamathecium inspersed; ascospores $20\text{--}25 \times 9\text{--}11 \mu\text{m}$; groups of fused ascomata dispersed to irregularly confluent; Singapore. *Astrothelium inspersogalbineum* Aptroot & Weerakoon
- J4b Hamathecium clear; ascospores variable in size. 5
- J5a Ascospores $21\text{--}28 \times 7\text{--}11 \mu\text{m}$. 6
- J5b Ascospores $28\text{--}55 \times 10\text{--}23 \mu\text{m}$. 7
- J6a Ascomata pseudostromatic; pseudostromata prominent, containing one or several groups of fused ascomata, covered with orange pigment; pantropical. *Astrothelium macrocarpum* (Fée) Aptroot & Lücking
- J6b Ascomata solitary to irregularly confluent, erumpent, containing a single group of fused ascomata; Thailand. *Astrothelium flavocoronatum* Luangsaph., Aptroot & Sangvichien
- J7a Ascospores $37\text{--}55 \times 15\text{--}23 \mu\text{m}$; ascomata pseudostromatic, pseudostromata erumpent, containing several groups of fused ascomata, covered with orange pigment; neotropical. *Astrothelium ochrothelium* (Nyl.) Müll. Arg.
- J7b Ascospores $28\text{--}35 \times 10\text{--}15 \mu\text{m}$. 8
- J8a Thallus strongly folded-bullate, whitish; groups of fused ascomata dispersed, immersed, with yellow-orange pigment around ostiolar area; pantropical. *Astrothelium versicolor* Müll. Arg.

- J8b Thallus smooth to uneven, light olive-brown to olive-green; ascomata pseudostromatic. 9
- J9a Pseudostromata immersed to erumpent, containing several groups of fused ascomata, forming irregular to reticulate lines covered with yellow-orange pigment; Brazil. *Astrothelium aureomaculatum* (Vain.) Aptroot & Lücking
- J9b Pseudostromata erumpent to prominent, each containing one group of fused ascomata, covered by orange pigment; New Caledonia. *Astrothelium grossum* Müll. Arg.
- J10a Ascospores 3-septate, rarely 3(–5)-septate. 11
- J10b Ascospores (3–)5–11-septate. 27
- J11a Ascospores small to medium-sized (20–52 × 6–17 µm). 12
- J11b Ascospores medium-sized to large (50–120 × 17–40 µm). 23
- J12a Ascospores 30–52 × 10–17 µm. 13
- J12b Ascospores 20–30 × 6–11 µm. 16
- J13a Hamathecium inspersion, inspersion dirty yellow-brown; lichexanthone on thallus and ascomata or ascomata only; ascomata pseudostromatic, pseudostromata prominent, containing one or several groups of fused ascomata. 14
- J13b Hamathecium clear; lichexanthone on thallus and ascomata; ascomata crowded or indistinctly pseudostromatic, pseudostromata immersed to erumpent, containing one or several groups of fused ascomata. 15
- J14a Lichexanthone on ascomata only; Brazil. *Astrothelium sordithecium* Lücking, M.P. Nelsen & Marcelli
- J14b Lichexanthone on thallus and ascomata; Brazil. *Astrothelium xanthosordithecium* Aptroot
- J15a Ascomata solitary or in dispersed, diffuse pseudostromata, conical with flattened top, whitish; neotropical. *Astrothelium leucothelium* Nyl.
- J15b Ascomata crowded, hemispherical to wart-shaped, covered by thallus; Malaysia. *Astrothelium chrysostomum* (Kremp.) Aptroot & Lücking
- J16a Lichexanthone on ascomata or only on ostiole. 17
- J16b Lichexanthone on thallus and ascomata; hamathecium inspersion or clear; ascospores 20–26 µm long. 21
- J17a Lichexanthone only on ostiole; hamathecium inspersion; Brazil. *Astrothelium eustominspersum* Aptroot & Oliveira-Junior

- J17b Lichexanthone on ascomata only; hamathecium clear or inspersed. 18
- J18a Hamathecium inspersed; Brazil. *Astrothelium studerae* Aptroot & M. Cáceres
J18b Hamathecium clear. 19
- J19a Hamathecium inspersed; ascomata diffusely pseudostromatic, erumpent, with whitish cover; Thailand. *Astrothelium neovariolosum* Luangsuph., Aptroot & Sangvichien
J19b Hamathecium clear. 20
- J20a Groups of fused ascomata dispersed, immersed to erumpent; Brazil. *Astrothelium laevigatum* Müll. Arg.
J20b Ascomata pseudostromatic, pseudostromata prominent, often whitish, containing one or several groups of fused ascomata; pantropical. *Astrothelium variolosum* (Ach.) Müll. Arg.
- J21a Ascomata diffusely pseudostromatic, pseudostromata erumpent, covered by thallus, containing one or several groups of fused ascomata; ascospores 20–25 µm; Brazil. *Astrothelium subinterjectum* Lücking, M.P. Nelsen & Jungbluth
J21b Ascomata distinctly pseudostromatic, pseudostromata prominent, with white or grey cover contrasting with the olive-green thallus, containing one or several groups of fused ascomata. 22
- J22a Thallus with pseudocyphellae; pseudostromata grey; Brazil. *Astrothelium pseudocyphellatum* R.C. Harris
J22b Thallus lacking pseudocyphellae; pseudostromata whitish; pantropical. *Astrothelium interjectum* R.C. Harris
- J23a Ascospores 100–130 × 35–45 µm; groups of fused ascomata dispersed to aggregate, immersed-erumpent, covered by thallus except for whitish ostiolar area; Venezuela. *Astrothelium ultralucens* Aptroot
J23b Ascospores 50–75 × 14–26 µm. 24
- J24a Lichexanthone on thallus (and sometimes also on ascomata). 25
J24a Lichexanthone not on thallus, only on ascomata. 26
- J25a Lichexanthone on thallus, not on ascomata; neotropical (Brazil and Venezuela). *Astrothelium leucoconicum* Nyl.
J25b Lichexanthone on thallus and ascomata; Bolivia. *Astrothelium nigrocacuminum* Flakus, Kukwa & Aptroot

- J26a Ascomata distinctly pseudostromatic, pseudostromata erumpent, with white cover contrasting with the olive-green thallus, containing one or several groups of fused ascomata; ascospores $60\text{--}70 \times 17\text{--}20 \mu\text{m}$; Brazil. *Astrothelium sinuosum* Aptroot & Gumboski
- J26b Groups of fused ascomata dispersed, immersed, with broad, dark ostiolar area; ascospores $50\text{--}60 \times 14\text{--}17 \mu\text{m}$; Brazil. *Astrothelium obtectum* Lücking, M.P. Nelsen & Benatti
- J27a Ascospores small ($20\text{--}30 \times 6\text{--}11 \mu\text{m}$), 3–5-septate. 28
- J27b Ascospores medium-sized to large ($30\text{--}110 \times 11\text{--}25 \mu\text{m}$), 5–11-septate. 30
- J28a Lichexanthone on ostioles only; ascomata distinctly pseudostromatic, pseudostromata erumpent, with white cover contrasting with the olive-green thallus, containing one or several groups of fused ascomata; pantropical. *Astrothelium eustomum* (Mont.) Müll. Arg.
- J28b Lichexanthone on thallus and ascomata; ascomata variable. 29
- J29a Ascomata distinctly pseudostromatic, pseudostromata erumpent, with white cover contrasting with the olive-green thallus; Thailand. *Astrothelium neglectum* Luangsaph., Aptroot & Sangvichien
- J29b Ascomata indistinctly pseudostromatic, pseudostromata erumpent to prominent, covered by thallus; pantropical. *Astrothelium sexloculatum* Aptroot
- J30a Ascospores $30\text{--}55 \mu\text{m}$ long; ascomata with white cover contrasting with the olive green thallus, erumpent. 31
- J30b Ascospores $65\text{--}110 \mu\text{m}$ long; ascomata variable. 34
- J31a Hamathecium inspersed; ascomata dispersed to irregularly confluent; Thailand. *Astrothelium siamense* Luangsaph., Aptroot & Sangvichien
- J31b Hamathecium clear; ascomata distinctly pseudostromatic, containing several groups of fused ascomata. 32
- J32a Lichexanthone on thallus and ascomata; neotropical (Guyana and Venezuela). *Astrothelium septemseptatum* Aptroot
- J32b Lichexanthone on ascomata or ostiole only. 33
- J33a Lichexanthone on ascomata; Brazil. *Astrothelium octoseptatum* Aptroot & M. Cáceres
- J33b Lichexanthone on ostiole only; French Guiana. *Astrothelium macroeustomum* Aptroot & Sipman

-
- J34a Hamathecium inspersed; lichexanthone on thallus and ascomata; ascomata distinctly pseudostromatic, pseudostromata erumpent, with white cover contrasting with the olive-green thallus, each containing a single group of fused ascomata; Cuba. *Astrothelium diplocarpoides* Müll. Arg.
- J34b Hamathecium clear; lichexanthone on ascomata only; ascomata variable. 35
- J35a Ascospores $80\text{--}110 \times 17\text{--}25 \mu\text{m}$, 5–7-septate; ascomata diffusely pseudostromatic, pseudostromata erumpent, covered by thallus, each containing a single group of fused ascomata; Brazil. *Astrothelium macrostomoides* Lücking, M.P. Nelsen & Benatti
- J35b Ascospores when transversely septate $65\text{--}85 \times 14\text{--}18 \mu\text{m}$, 5–11-septate, submuriform when longer; ascomata variable. 36
- J36a Ascospores 5–7-septate; ascomata diffusely pseudostromatic, pseudostromata erumpent to prominent, covered by thallus, uneven-verrucose, each containing a single group of fused ascomata; French Guiana. *Astrothelium macrostomum* Aptroot
- J36b Ascospores 9–11-septate; ascomata distinctly pseudostromatic, pseudostromata immersed, with white cover contrasting with the olive-green thallus, containing one to several groups of fused ascomata. 37
- J37a Ascospores $65\text{--}70 \times 15\text{--}17 \mu\text{m}$, transversely septate only; Brazil. *Astrothelium eumultiseptatum* Aptroot & M. Cáceres
- J38b Ascospores $90\text{--}110 \times 22\text{--}28 \mu\text{m}$, in part with longitudinal septa and becoming submuriform; neotropical. *Astrothelium diplocarpum* Nyl.
- J38a Ascospores 5–9-septate. 39
- J38b Ascospores 3-septate. 41
- J39a Ascospores 5-septate; ascomata diffusely pseudostromatic, pseudostromata with external pink pigment; Brazil. *Astrothelium bivelum* Aptroot & M. Cáceres
- J39b Ascospores 5–9-septate. 40
- J40a Ascospores 7–9-septate; only ostiole yellow pruinose; Brazil. *Astrothelium septoconicum* Aptroot
- J40b Ascospores 6–7-septate; pseudostromata with internal yellow pigment; Brazil. *Astrothelium introflavidum* Aptroot & M. Cáceres
- J41a Ascospores small ($20\text{--}30 \times 6\text{--}12 \mu\text{m}$). 42
- J41b Ascospores small to medium-sized ($30\text{--}65 \times 10\text{--}25 \mu\text{m}$). 47
- J42a Thallus and ascomata with orange pigment. 43

- J42b Pigment on ascomata only, yellow to orange. 44
- J43a Pseudostromata immersed to erumpent, whitish; Indonesia. *Astrothelium pseudoferrugineum* Aptroot
- J43b Pseudostromata erumpent to prominent, covered by thallus; Cuba. *Astrothelium ferrugineum* (Müll. Arg.) Aptroot & Lücking
- J44a Hamathecium inspersed; Sri Lanka. *Astrothelium inspersoconicum* Aptroot & Weerakoon
- J44b Hamathecium clear. 45
- J45a Pigment yellow, sparse; ostioles partly apical; ascomata solitary to irregularly confluent and fused, erumpent, covered by thallus; Bolivia. *Astrothelium pallidoflavum* Flakus & Aptroot
- J45b Pigment orange, distinct; ostioles consistently lateral and fused; ascomata pseudostromatic, erumpent to prominent, with orange cover, containing one to many of fused ascomata. 46
- J46a Ascomata with one or few fused ascomata; pantropical. *Astrothelium cinnamomeum* (Eschw.) Müll. Arg.
- J46b Ascomata pseudostromatic, with many groups of fused ascomata; Brazil. *Astrothelium stromatocinnamomeum* Aptroot
- J47a Thallus yellow; ascospores $30\text{--}35 \times 10\text{--}12 \mu\text{m}$; Brazil. *Astrothelium croceum* Malme
- J47b Thallus not cracked; ascospores $(30\text{--})35\text{--}65 \times (10\text{--})12\text{--}25 \mu\text{m}$. 48
- J48a Ascospores $55\text{--}65 \times 15\text{--}25 \mu\text{m}$; pantropical. *Astrothelium ocellatum* Malme
- J48b Ascospores $30\text{--}45 \times 10\text{--}17 \mu\text{m}$; pantropical. *Astrothelium scorioides* Nyl.

Key K. *Astrothelium* species with ascospores transversely septate, ostiole lateral, secondary substances absent

- K1a Ascospores 3-septate. 2
- K1b Ascospores (3–)5–23-septate. 21
- K2a Ascospores medium-sized ($55\text{--}80 \times 15\text{--}25 \mu\text{m}$); thallus distinctly verrucose papillose. 3
- K2b Ascospores small to medium-sized ($(12\text{--})20\text{--}45 \times (4\text{--})7\text{--}15 \mu\text{m}$); thallus variable. 4

- K3a Ascospores medium-sized (55–65 × 15–19 µm); thallus distinctly verrucose-papillose; ascomata immersed; Brazil. *Astrothelium simplex* Aptroot & S.M.A. Martins
- K3b Ascospores 80–88 × 32–36 µm; ascomata prominent to sessile; Panama. *Astrothelium pseudannulare* Aptroot & Etayo
- K4a Hamathecium inspersed. 5
- K4b Hamathecium clear. 7
- K5a Thallus covering the pseudostromata; Sri Lanka. *Astrothelium conjugatum* Weerakoon & Aptroot
- K5b Thallus not covering the pseudostromata. 6
- K6a Ascospores 20–25 µm long; ascomata diffusely pseudostromatic, pseudostromata prominent, laterally covered by thallus, with one to several groups of fused ascomata forming broad, flat, dark ostiolar areas often fused in lobate pattern; eastern palaeotropical. *Astrothelium straminicolor* (Nyl.) Aptroot & Lücking
- K6b Ascospores 25–30 µm long; ascomata diffusely pseudostromatic, pseudostromata prominent, covered by thallus, with one to several groups of fused ascomata with dark, papilliform, always separate ostiolar areas; pantropical. *Astrothelium pyrenastrosulphureum* Aptroot & Lücking
- K7a Ascospores (12–)20–27 × (4–)7–10 µm. 8
- K7b Ascospores 28–45 × 10–15 µm. 17
- K8a Ascospores 12–15 × 4–5 µm; ascomata diffusely pseudostromatic, pseudostromata prominent, laterally covered by thallus, with one to several groups of fused ascomata forming broad, flat, dark ostiolar areas often fused in lobate pattern; Cuba. *Astrothelium acrophaeum* Müll. Arg.
- K8b Ascospores 19–27 × 7–10 µm; thallus and ascomata variable (*Astrothelium crassum* complex). 9
- K9a Ostioles separate; ascomata aggregate in irregular groups or lines, immersed; neotropical. *Astrothelium scorizum* (Müll. Arg.) Aptroot & Lücking
- K9b Ostioles fused; fused ascomata dispersed to aggregate. 10
- K10a Thallus mottled olive-whitish, with irregular whitish areas or lines; ascomata usually developed on the whitish areas (but not covered by them). 11
- K10b Thallus uniformly olive-brown to olive-green, without whitish areas except when covering the ascomata. 13

- K11a Ascomata dispersed, conical, partially or fully covered by thallus; Malaysia.
Astrothelium peranceps (Kremp.) Aptroot & Lücking
- K11b Ascomata dispersed to irregularly confluent, completely exposed. 12
- K12a Ascomata black, developed on distinct, white, irregular to reticulate lines; Brazil.
Astrothelium nigricans Malme
- K12b Ascomata dark brown, emerging from thallus; neotropical. *Astrothelium nigrum*
Aptroot & M. Cáceres
- K13a Ascomata aggregate in irregular groups bordered by a thin whitish line; thallus
rather thick, uneven to irregularly folded; eastern palaeotropical. *Astrothelium*
subfuscum Kremp.
- K13b Ascomata dispersed to irregularly confluent, exposed and brown or with whitish or
(partial) thallus cover; thallus thin to thick but not folded. 14
- K14a Ascomata exposed and brown-black, conical; Brazil. *Astrothelium obscurum* Müll.
Arg.
- K14b Ascomata at least with lateral thallus cover or with whitish rim or cover, lens-
shaped to hemispherical. 15
- K15a Ascomata prominent, with basal thallus cover and upper part exposed; individual
ascomata clearly discernible; Australia. *Astrothelium oligocarpum* (Müll. Arg.)
Aptroot & Lücking
- K15b Ascomata erumpent to prominent, whitish or whitish rim; individual ascomata only
visible as immersed chambers. 16
- K16a Ascomata with complete whitish cover including ostiolar area; thallus thin;
Thailand. *Astrothelium subdissocians* (Nyl. ex Vain.) Aptroot & Lücking
- K16b Ascomata with whitish rim bordering a wide, dark ostiolar area; thallus thickened;
pantropical. *Astrothelium crassum* (Fée) Aptroot
- K17a Thallus strongly verrucose-bullate; groups of fused ascomata dispersed, immersed,
visible only by their dark, narrow ostiolar area; Brazil. *Astrothelium globosum*
Aptroot & M. Cáceres
- K17b Thallus smooth to uneven; ascomata distinct, diffusely to distinctly
pseudostromatic, pseudostromata covered with thallus. 18
- K18a Ascomata distinctly pseudostromatic, pseudostromata strongly prominent to sessile,
uneven-verrucose, covered with thallus, with whitish ring around dark ostiolar
zone, containing one or several groups of fused ascomata; pantropical.
Astrothelium fallax Müll. Arg.

- K18b Ascomata diffusely pseudostromatic, pseudostromata erumpent to prominent, smooth to uneven, covered with thallus, containing several to many groups of fused ascomata. 19
- K19a Ascospores $35\text{--}45 \times 14\text{--}16 \mu\text{m}$; Brazil. *Astrothelium marcidum* (Fée) Aptroot & Lücking
- K19b Ascospores $28\text{--}33 \times 8\text{--}15 \mu\text{m}$. 20
- K20a Pseudostromata with distinctly separate, whitish ostiolar areas; ascospores $12\text{--}15 \mu\text{m}$ broad; Costa Rica. *Astrothelium intermedium* Aptroot & Lücking
- K20b Pseudostromata with upper part entirely whitish; ascospores $8\text{--}10 \mu\text{m}$ broad; Brazil. *Astrothelium pleiostomum* Redinger
- K21a Ascospores small and slender, 6–9-septate, $35\text{--}54 \times 7\text{--}10 \mu\text{m}$, curved; India. *Astrothelium himalayense* (Upreti & Aptroot) Aptroot & Schumm
- K21b Ascospores medium-sized ($45\text{--}80 \times 12\text{--}28 \mu\text{m}$). 22
- K21c Ascospores large ($80\text{--}130 \times 17\text{--}30 \mu\text{m}$). 29
- K22a Hamathecium inspersed. 23
- K22b Hamathecium clear. 24
- K23a Ascospores $45\text{--}55 \times 13\text{--}16 \mu\text{m}$, 7–9-septate; ascomata indistinctly pseudostromatic, pseudostromata erumpent to prominent, brownish; Venezuela. *Astrothelium tryptelioides* Aptroot
- K23b Ascospores $50\text{--}75 \times 14\text{--}22 \mu\text{m}$, 5-septate; ascomata diffusely pseudostromatic, pseudostromata erumpent, covered by thallus, containing one to several groups of fused ascomata; pantropical. *Astrothelium subclandestinum* Leight.
- K23c Ascomata in distinct, white pseudostromata; Vietnam. *Astrothelium decemloculare* Aptroot, Weerakoon & G. Vo
- K24a Ascospores 4–5-septate; groups of fused ascomata dispersed, erumpent, with whitish ring around dark ostiolar zone. 25
- K24b Ascospores 7–19-septate; ascomata variable. 26
- K25a Ascospores predominantly 4-septate, $45\text{--}55 \times 15\text{--}18 \mu\text{m}$; Brazil. *Astrothelium quatuorseptatum* Aptroot & M. Cáceres
- K25b Ascospores 5-septate, $50\text{--}75 \times 14\text{--}22 \mu\text{m}$; Brazil. *Astrothelium supraclandestinum* Aptroot & M. Cáceres
- K26a Ascospores 7-septate, $14\text{--}28 \mu\text{m}$ broad; ascomata pseudostromatic, pseudostromata immersed-erumpent, with whitish cover; neotropical (Brazil and Guyana). *Astrothelium zebrinum* Aptroot

- K26b Ascospores 9–19-septate, 12–17 μm broad. 27
- K27a Ascospores 45–55 μm long, consistently 9-septate; ascomata pseudostromatic, pseudostromata erumpent, with white cover contrasting with olive-green thallus, with several to many groups of fused ascomata; Brazil. *Astrothelium novemseptatum* Aptroot & M. Cáceres
- K27b Ascospores 60–80 μm long, 9–19-septate. 28
- K28a Ascospores 69–80 μm long, 11–15-septate; ascomata deeply immersed in groups; Brazil. *Astrothelium valsoides* Aptroot & M. Cáceres
- K28b Ascospores 60–80 μm long, 9–19-septate; groups of fused ascomata dispersed, erumpent, brown-black with whitish ring around base; eastern palaeotropical. *Astrothelium heterophorum* Nyl.
- K29a Ascospores I+ violet, occasionally with a longitudinal septum in one or several cells; thallus bullate-folded; groups of ascomata dispersed to irregularly confluent, immersed to erumpent, exposed and brown-black; Bolivia. *Astrothelium neodiplocarpum* Flakus, Kukwa & Aptroot
- K29b Ascospores I–, lacking longitudinal septa; groups of fused ascomata dispersed (or ostioles separate), mostly covered by thallus. 30
- K30a Hamathecium inspersed; groups of fused ascomata prominent, with broad white ostiolar area; ascospores 9–11-septate, 80–100 μm long; Thailand. *Astrothelium macrostiolum* Luangsaph., Aptroot & Sangvichien
- K30b Hamathecium clear; ascomata variable; ascospore septa variable, ascospores (90–)100–130 μm long. 31
- K31a Ascospores 5–7(–9)-septate. 32
- K31b Ascospores 9–23-septate. 33
- K32a Ascospore median septum not thickened; neotropical. *Astrothelium robustum* Müll. Arg.
- K32b Ascospore median septum thickened; Brazil. *Astrothelium fernandae* Aptroot
- K33b Ascospores 9–11-septate, median septum thickened; ostioles fused; Brazil. *Astrothelium medioincrassatum* Aptroot & M. Cáceres
- K33c Ascospores 11–23-septate. 34
- K34a Ascospores 11–15-septate; groups of fused ascomata erumpent; Brazil. *Astrothelium robustosporum* Aptroot & M. Cáceres

K34b Ascospores 19–23-septate; groups of fused ascomata strongly prominent to sessile, with protruding, irregular ostioles; neotropical. *Astrothelium gigasporum* R.C. Harris

Key L. *Astrothelium* species with ascospores muriform, ostiole apical, secondary substances present on thallus and/or ascoma

L1a Lichexanthone present; thallus and/or ascomata UV+ yellow; pigment(s) present or absent. 2

L1b Lichexanthone absent; pigment(s) always present, thallus and ascomata usually K+ red to purple and UV+ orange or red. 14

L2a Pigments present on/in thallus and/or ascomata; ascospores large, 70–160 × 15–32 µm. 3

L2b Pigments absent; ascospores medium-sized to large, 40–160 × 15–40 µm. 6

L3c Hamathecium inspersed; ascomata mostly solitary; lichexanthone only on thallus. Vietnam. *Astrothelium curcumarum* Aptroot, Weerakoon & G. Vo

L3b Hamathecium clear. 4

L4a Thallus and ascomata with external, yellow-orange, K+ purple pigment(s); lichexanthone on thallus only; ascomata distinctly pseudostromatic, pseudostromata erumpent to prominent, covered with orange pigment, sharply delimited and contrasting with thallus; Brazil. *Astrothelium flavoduplex* Aptroot & M. Cáceres

L4b Ascomata or medulla with internal, red, K+ yellow-green pigment(s); lichexanthone on thallus and ascomata. 5

L5a Ascospores I+ violet; ascomata solitary to irregularly confluent, erumpent to prominent, whitish around ostiole(s); Brazil. *Astrothelium sanguineoxanthum* Aptroot

L5b Ascospores I–; ascomata distinctly pseudostromatic, pseudostromata erumpent to prominent, whitish, sharply delimited and contrasting with thallus; Bolivia. *Astrothelium elixii* Flakus & Aptroot

L6a Hamathecium inspersed. 7

L6b Hamathecium clear. 10

L7a Ascospores (2–)4(–6) per ascus, 40–80 µm long; on rock; Brazil. *Astrothelium stramineum* (Malme) Aptroot & Lücking

L7b Ascospores 8 per ascus. 8

- L8a Ascospores 65–71 μm long; Brazil. *Astrothelium rhinohallinum* Kalb & Aptroot
L8b Ascospores 70–220 μm long. 9
- L9a Ascospores 70–120 \times 18–24 μm ; eastern palaeotropical. *Astrothelium meristosporoides* (P.M. McCarthy & Vongsheew.) Aptroot & Lücking
L9b Ascospores 120–220 \times 25–40 μm ; eastern palaeotropical. *Astrothelium meristosporum* (Mont. & Bosch) Aptroot & Lücking
- L10a Ascospores 70–115 \times 15–24 μm . 11
L10b Ascospores 110–170 \times 20–35 μm . 13
- L11a Lichexanthone on pseudostromata only; ascomata covered by thallus except for dark ostiolar area; neotropical (Brazil and Guyana). *Astrothelium corallinum* Aptroot
L11b lichexanthone on thallus, sometimes also on ascomata. 12
- L12a Ascospores 70–90 \times 18–20 μm , 8 per ascus; lichexanthone on thallus only; ascomata indistinctly pseudostromatic, pseudostromata erumpent, whitish, not sharply delimited; Guyana. *Astrothelium lucidohallinum* Aptroot
L12b Ascospores 80–115 \times 15–24 μm , 4 per ascus; lichexanthone on thallus and ascomata; ascomata distinctly pseudostromatic, pseudostromata erumpent to prominent, brown-black, sharply delimited; Brazil. *Astrothelium ochroleucooides* Aptroot & M. Cáceres
- L13a Ascomata pseudostromatic; pseudostromata erumpent, whitish; Bolivia. *Astrothelium variabile* Flakus & Aptroot
L13b Ascomata pseudostromatic; pseudostromata erumpent, brown-black; Brazil. *Astrothelium xanthosuperbum* Aptroot & M. Cáceres
- L14a Ascospores medium-sized, 45–80 μm long. 15
L14b Ascospores large over 80 μm long. 21
- L15a Ascospores 40–55 μm long. 16
L15b Ascospores 60–80 μm long. 18
- L16a Ascomata with internal, yellow pigment; ascospores about 3–6 times as long as broad. 17
L16b Thallus and ascomata with external, orange pigment and medulla (ascomata) additionally with internal, purple pigment; ascospores about twice as long as broad; Ecuador. *Astrothelium condoricum* Aptroot

-
- L17a Ascospores about 4–6 times as long as broad; Brazil. *Astrothelium lineatum* Aptroot & M. Cáceres
- L17b Ascospores about 3–4 times as long as broad; Brazil. *Astrothelium duplicatum* Aptroot & M. Cáceres
- L18a Ascomata with internal red pigment crystals; ostiolar area very broad and flat, black; Sierra Leone. *Astrothelium sierraleonense* (C.W. Dodge) Aptroot & Lücking
- L18b Pigment external; ostiolar area narrow, whitish. 19
- L19a Hamathecium inspersed; pigment on ascomata only; ascospores 20–30 µm broad; ascomata pseudostromatic, pseudostromata prominent, covered with orange pigment, sharply delimited from thallus; neotropical. *Astrothelium auratum* (R.C. Harris) Aptroot & Lücking
- L19b Hamathecium clear; pigment on thallus and ascomata; ascospores 12–25 µm broad; ascomata variable. 20
- L20a Ascomata pseudostromatic; pseudostromata erumpent, covered with orange pigment, sharply delimited; Brazil. *Astrothelium graphicum* Aptroot & S.M.A. Martins
- L20b Ascomata pseudostromatic; pseudostromata immersed, whitish, not sharply delimited; neotropical. *Astrothelium flavomaculatum* Aptroot
- L21a Ascomata with internal, red, K+ yellow-green pigment; ascospores I + violet; ascomata pseudostromatic, pseudostromata erumpent to prominent, brown-black, sharply delimited; Brazil. *Astrothelium sanguinarium* (Malme) Aptroot & Lücking
- L21b Thallus and/or ascomata with external or internal, yellow to orange, K+ purple pigment; ascospores I–; ascomata variable. 22
- L22a Pigment in medulla of pseudostromata only. 23
- L22b Pigment external on thallus and/or ascomata. 25
- L23a. Ascospores 8 per ascus, 90–105 × 26–36 µm; Malaysia. *Astrothelium deforme* (Fée) Aptroot & Lücking
- L23b Ascospores 2 or 4 per ascus. 24
- L24a Ascospores 2 per ascus, 110–130 × 30–35 µm; Guyana. *Astrothelium laurerosphaerioides* Aptroot
- L24a Ascospores 4 per ascus, up to 330 × 40 µm; French Guiana. *Astrothelium cayennense* Aptroot & Sipman

L25a Ascospores 8 per ascus, 20–25 µm broad; ascomata pseudostromatic, immersed to erumpent, covered with orange pigment, sharply delimited; Brazil. *Astrothelium mesoduplex* Aptroot & M. Cáceres

L25b Ascospores 2–4 per ascus, 25–50 µm broad; ascomata variable. 26

L26a Pigment on ascomata only, absent from thallus; ostiole in part lateral; Caribbean. *Astrothelium chrysoglyphum* (Vain.) Aptroot & Lücking

L26b Pigment on thallus only, absent from ascomata; ostiole consistently apical; pantropical. *Astrothelium vezdae* (Makhija & Patw.) Aptroot & Lücking

Key M. *Astrothelium* species with ascospores muriform, ostiole apical, secondary substances absent

M1a Hamathecium inspersed. 2

M1b Hamathecium clear. 20

M2a Ascospores small to medium-sized (30–60 × 11–20 µm). 3

M2b Ascospores large (80–300 × 20–75 µm). 6

M3a Ascomata aggregate into irregular groups or lines, immersed to erumpent, exposed, blackish; ascospores 35–50 × 12–18 µm; pantropical. *Astrothelium subdiscretum* (Nyl.) Aptroot & Lücking

M3b Ascomata solitary or few aggregate, at least partly covered by thallus; ascospores variable. 4

M4a Ascospores 32–45 × 11–13 µm; ascomata aggregate to pseudostromatic, pseudostromata prominent to sessile, covered by thallus; India. *Astrothelium pseudovariatum* (Upreti & Ajay Singh) Aptroot & Lücking

M4b Ascospores 45–70 × 13–20 µm; ascomata solitary to aggregate, if aggregate then whitish. 5

M5a Ascomata solitary, prominent, mostly exposed and brown-black; Brazil. *Astrothelium ambiguum* (Malme) Aptroot & Lücking

M5b Ascomata solitary, erumpent, mostly covered by thallus except for dark ostiolar area; eastern palaeotropical. *Astrothelium indicum* (Upreti & Ajay Singh) Aptroot & Lücking

M6a Ascospores 1–2 per ascus, 210–300 µm long. 7

M6b Ascospores 4–8 per ascus, 80–200(–230) µm long. 8

M6c Ascospores 4 per ascus, 240–260 µm long; Brazil. *Astrothelium flavogigasporum* Aptroot

- M7a Ascospores 1 per ascus, 240–300 × 45–50 µm, with thickened median septum; ascomata confluent to irregularly pseudostromatic, immersed-erumpent, with whitish cover around the blackish ostiole; Colombia. *Astrothelium colombiense* Aptroot
- M7b Ascospores 2 per ascus, 210–225 × 30–35 µm, without thickened median septum; ascomata solitary, prominent, mostly exposed and brown-black; eastern palaeotropical. *Astrothelium megaleium* (Kremp.) Aptroot & Lücking
- M8a Ascospores submuriform, with few longitudinal septa, 17–22 µm broad; ascomata solitary to irregularly confluent, immersed to erumpent, covered by thallus with broad whitish area around the dark ostiolar spot; India. *Astrothelium macrosporum* (Makhija & Patw.) Aptroot & Lücking
- M8b Ascospores distinctly muriform, 20–75 µm broad; ascomata variable. 9
- M9a Ascospores 80–140 µm long. 10
- M9b Ascospores (125–)140–230 µm long, most ascospores distinctly exceeding 140 µm. 12
- M10a Ascospores 2–3 times as long as broad (80–100(–125) × 28–48 µm), without thickened median septum; ascomata solitary, erumpent, laterally covered by thallus with ostiolar area exposed, black; neotropical. *Astrothelium effusum* (Aptroot & Sipman) Aptroot & Lücking
- M10b Ascospores 4–5 times as long as broad (115–140 × 25–33 µm), with thickened median septum; ascomata variable. 11
- M11a Ascomata solitary, strongly prominent, completely covered by thallus; eastern palaeotropical. *Astrothelium aurantiacum* (Makhija & Patw.) Aptroot & Lücking
- M11b Ascomata solitary to irregularly confluent, immersed to erumpent, covered by thallus but with exposed, dark ostiolar area bordered by whitish rim; eastern palaeotropical. *Astrothelium sikkimense* (Makhija & Patw.) Aptroot & Lücking
- M12a Ascospores with thickened median septum. 13
- M12b Ascospores without thickened median septum. 17
- M13a Ascomata distinctly pseudostromatic, pseudostromata prominent to sessile, with whitish cover; thallus smooth to uneven; Indonesia. *Astrothelium alboverrucoides* Aptroot
- M13b Ascomata solitary; thallus variable. 14
- M14a Thallus smooth to uneven; ascomata prominent, exposed and blackish; pantropical. *Astrothelium flavomeristosporum* Aptroot

- M14b Thallus verrucose-papillose or bullate; ascomata immersed, covered by thallus. 15
- M15a Thallus coarsely bullate, gall-like; ascomata immersed in large thallus verrucae, completely covered by thallus; Philippines. *Astrothelium philippinense* Aptroot & Schumm
- M15b Thallus verrucose-papillose; ascomata variable but ostioles partly lateral. 16
- M16a Hamathecium orange; ascospores 26–34 µm broad; Papua New Guinea. *Astrothelium papillosum* (P.M. McCarthy) Aptroot & Lücking
- M16b Hamathecium uncoloured; ascospores 35–45 µm broad; Venezuela. *Astrothelium komposchii* Aptroot
- M17a Ascospores 135–165 µm long; thallus uneven-verrucose; ascomata solitary to irregularly confluent, erumpent, covered by thallus, uneven-verrucose; Brazil. *Astrothelium irregulare* (Müll. Arg.) Aptroot & Lücking
- M17b Ascospores (140–)165–230 µm long; thallus and ascomata variable. 18
- M18a Hamathecium yellow; thallus bullate; ascomata solitary, prominent, beaked; neotropical. *Astrothelium flavomegaspermum* Aptroot & Etayo
- M18b Hamathecium hyaline. 19
- M19a Thallus bullate; ascomata solitary to confluent, immersed in coarse thallus verrucae, covered by thallus, with yellowish ostiolar rim; Ecuador. *Astrothelium flavostiolum* Aptroot
- M19b Thallus smooth to uneven; ascomata solitary, prominent, covered by thallus except for dark ostiolar area usually bordered by a whitish rim; pantropical. *Astrothelium megaspermum* (Mont.) Aptroot & Lücking
- M20a Ascospores small (30–60 × 11–20 µm). 21
- M20b Ascospores medium-sized to large (60–280 × (15–)20–65 µm). 22
- M21a Ascospores 30–45 × 11–13 µm; ascomata pseudostromatic, pseudostromata (strongly) prominent, covered by thallus except for ostioles; neotropical. *Astrothelium variatum* (Nyl.) Aptroot & Lücking
- M21b Ascospores 50–60 × 15–20 µm; ascomata solitary, prominent, covered by thallus; eastern palaeotropical. *Astrothelium keralense* (Upreti & Ajay Singh) Aptroot & Lücking
- M22a Ascospores 200–280 × 50–65 µm, 2 per ascus; ascomata irregularly grouped to diffusely pseudostromatic, pseudostromata erumpent, completely covered by thallus; neotropical. *Astrothelium gigantosporum* (Müll. Arg.) Aptroot & Lücking
- M22b Ascospores 70–195 × 14–45 µm, (2–)4–8 per ascus; ascomata variable. 23

- M23a Ascospores I+ blue. 24
M23b Ascospores I-negative. 25
- M24a Ascospores I+ violet-blue, 125–170 × 27–40 µm; Bolivia. *Astrothelium amylosporum* Flakus & Aptroot
M24b Ascospores I+ pale blue, 85–100 × 20–24 µm; Thailand. *Astrothelium palaeoexostemmatidis* Sipman & Aptroot
- M25a Ascospores with thickened median septum. 26
M25b Ascospores without thickened median septum. 27
- M26a Ascospores 4 per ascus, 145–175 × 30–35 µm; thallus strongly bullate; ascomata solitary to irregularly confluent, immersed in bullate thallus warts and completely covered by thallus except for protruding ostiole; Brazil. *Astrothelium tetrasporum* Aptroot & M. Cáceres
M26b Ascospores (6–)8 per ascus, (85–)100–170 × 23–33 µm; thallus smooth to uneven; ascomata solitary to diffusely pseudostromatic, prominent, whitish surrounding the black ostiolar area; India. *Astrothelium alboverrucum* (Makhija & Patw.) Aptroot & Lücking
- M27a Ascospores 1–4 per ascus. 28
M27b Ascospores 8 per ascus. 31
- M28a Ascospores 1 per ascus, 125–148 µm long; Brazil. *Astrothelium unisporum* Aptroot & M. Cáceres
M28b Ascospores 1–4 per ascus. 29
- M29a Ascomata distinctly pseudostromatic, pseudostromata erumpent to prominent, with white cover, sharply delimited from thallus; ascospores 80–140 × 25–35 µm; pantropical. *Astrothelium sphaerioides* (Müll. Arg.) Aptroot & Lücking
M29b Ascomata irregularly confluent to diffusely pseudostromatic, erumpent to prominent, either fully exposed and brown-black or covered by olive-green thallus; ascospores 120–180(–200) × 30–50(–60) µm. 30
- M30a Ascomata erumpent, fully exposed and brown-black; ascospores 120–175 × 30–45 µm; neotropical. *Astrothelium subdisjunctum* (Mont.) Aptroot & Lücking
M30b Ascomata prominent, covered by thallus except dark ostiole bordered by a white rim, with one to several ascomata per thallus wart and ostioles often pointing in different directions; ascospores 160–180(–200) × 40–50(–60) µm; Central America. *Astrothelium nicaraguense* Lücking, M.P. Nelsen & T. Orozco

- M31a Ascospores $150\text{--}190 \times 40\text{--}45 \mu\text{m}$; ascomata solitary or few grouped, covered by thallus; neotropical (Brazil and Bolivia). *Astrothelium octosporoides* Aptroot & Lücking
- M31b Ascospores $60\text{--}150 \times 15\text{--}30 \mu\text{m}$; ascomata variable. 32
- M32a Ascomata solitary, immersed to erumpent, covered by thallus except for dark ostiolar area surrounded by irregular whitish rim; ascospores not exceeding $80 \mu\text{m}$ in length; neotropical. *Astrothelium exostemmatitis* (Müll. Arg.) Aptroot & Lücking
- M32b Ascomata irregularly confluent to pseudostromatic, erumpent to prominent, fully exposed; ascospores usually exceeding $80 \mu\text{m}$ in length; Brazil. *Astrothelium chapadense* (Malme) Aptroot & Lücking

Key N. *Astrothelium* species with ascospores muriform, ostiole lateral, secondary substances present on thallus and/or ascoma

- N1a Lichexanthone present; thallus and/or ascomata UV+ yellow; pigment(s) present or absent. 2
- N1b Lichexanthone absent; pigment(s) always present, thallus and ascomata usually K+ red to purple and UV+ orange or red. 17
- N2a Hamathecium inspersed; ascomata with separate or fused ostioles. 3
- N2b Hamathecium clear; ascomata with fused ostioles. 6
- N3a Ascospores $80\text{--}115 \times 25\text{--}35 \mu\text{m}$, 4 per ascus; lichexanthone in medulla; ascomata with separate ostioles, immersed, covered by thallus; Ecuador. *Astrothelium lucidomedullatum* Aptroot
- N3b Ascospores $150\text{--}220 \times 40\text{--}65 \mu\text{m}$, 2 or 8 per ascus; lichexanthone external on thallus and/or ascomata. 4
- N4a Ascospores 8 per ascus, $40\text{--}50 \mu\text{m}$ broad; ascomata with separate ostioles,; lichexanthone on thallus and ascomata. 5
- N4b Ascospores 2 per ascus, $50\text{--}65 \mu\text{m}$ broad; ascomata with fused ostioles, groups of fused ascomata dispersed, covered by thallus except ostioles; lichexanthone on ascomata only; Brazil. *Astrothelium megalostomum* (Vain.) Aptroot & Lücking
- N5a Ascomata covered by thallus except for broad black ostiolar area; Bolivia. *Astrothelium carrascoense* Flakus, Kukwa & Aptroot
- N5b Ascomata with separate ostioles, covered by thallus, without black ostiolar area; Brazil. *Astrothelium aureoirregulare* Aptroot & Gumboski
- N6a Ascospores small ($35\text{--}45 \times 15\text{--}19 \mu\text{m}$). 7

- N6b Ascospores large (85–250 × 19–65 µm). 8
- N7a Only ostiole UV+ yellow; fused ascomata pseudostromatic, pseudostromata erumpent, with white cover contrasting with olive-green thallus; Brazil.
Astrothelium eustomurale Aptroot & M. Cáceres
- N7b Entire pseudostroma UV+ yellow; fused ascomata pseudostromatic, pseudostroma immersed, with white cover contrasting with olive-brown thallus; Guyana.
Astrothelium lucidostromum Aptroot
- N8a Ascospores 150–250 × 30–65 µm, (1–)2 per ascus; ascomata with fused ostioles. 9
N8b Ascospores 85–150 × 19–45 µm, 2 or 8 per ascus; ascomata variable. 12
- N9a Fused ascomata solitary, immersed and covered by thallus; lichexanthone on thallus and ascomata; neotropical. *Astrothelium sepultum* (Mont.) Aptroot & Lücking
- N9b Fused ascomata pseudostromatic, pseudostromata immersed, whitish and contrasting with olive-green thallus; thallus bullate-bumpy; lichexanthone on pseudostromata only. 10
- N10a Lichexanthone only on the thallus; pseudostroma with orange pigment; Brazil.
Astrothelium astrolucidum Aptroot & M. Cáceres
- N10b Lichexanthone only on pseudostromata; pseudostromata without pigment. 11
- N11a Thallus dominated by finely and densely cracked pseudostromata, with thallus bumps dispersed; ascomata completely immersed in pseudostromata, hardly visible; Panama. *Astrothelium cryptolucens*
Lücking, M.P. Nelsen & N. Salazar
- N11b Thallus dominated by vegetative part forming large bumps separating the smooth pseudostromata; ascomata immersed to erumpent with the ostiolar area broadly exposed and black; neotropical. *Astrothelium norisianum* Lücking, M.P. Nelsen & Aptroot
- N12a Ascospores 85–130 µm long, 2 per ascus; lichexanthone internal in medulla; fused ascomata solitary to irregularly confluent, immersed to erumpent, covered by thallus except dark ostiolar area surrounded by irregular whitish rim; neotropical.
Astrothelium consimile (Müll. Arg.) Aptroot & Lücking
- N12b Ascospores 100–150 µm long, 8 per ascus; lichexanthone external on ascomata; ascomata variable. 13
- N13a Ascospores 30–45 µm broad. 14
N13b Ascospores 17–23 µm broad; ascomata variable. 15

- N14a Ascomata solitary; lichexanthone on pseudostromata; neotropical. *Astrothelium defossum* (Müll. Arg.) Aptroot & Lücking
- N14b Ascomata fused; lichexanthone only on ostiole; Brazil. *Astrothelium megeustomurale* M. Cáceres & Aptroot
- N15a Ascomata solitary, immersed to erumpent, covered by thallus except dark ostiolar area; thallus shallowly folded; Brazil. *Astrothelium octosporum* (Vain.) Aptroot & Lücking
- N15b Ascomata irregularly confluent to grouped, erumpent to prominent, completely covered by thallus; thallus smooth to uneven. 16
- N16a Ostioles visible as sessile, broadly white papillae; Brazil. *Astrothelium megeustomum* Aptroot & Fraga Jr.
- N16b Ostioles visible as minute, immersed dark pores; neotropical. *Astrothelium confluens* (Müll. Arg.) Aptroot & Lücking
- N17a Ascospores (small to) medium-sized ($35\text{--}80 \times 12\text{--}25 \mu\text{m}$). 18
- N17b Ascospores large ($75\text{--}200 \times 22\text{--}50 \mu\text{m}$). 20
- N18a pigment external on ascomata only, orange. 19
- N18b Ascomata pseudostromatic, pseudostromata immersed, whitish, forming irregular to reticulate lines, well delimited from thallus; pigment external on thallus only, orange; French Guiana. *Astrothelium praetervisum* (Müll. Arg.) Aptroot & Lücking
- N19a Ascomata pseudostromatic, with fused ostioles; Brazil. *Astrothelium muriconicum* Aptroot
- N19b Ascomata solitary, immersed to erumpent, covered by thallus except dark ostiolar area surrounded by whitish rim; neotropical. *Astrothelium subaequans* Müll. Arg.
- N20a Pigment orange, external on ascomata or internal in medulla. 21
- N20b Pigment red, external on ascomata. 23
- N21a Pigment internal in medulla of ascomata; neotropical. *Astrothelium amazonum* (R.C. Harris) Aptroot & Lücking
- N21b Pigment external on ascomata. 22
- N22a Ascospores $100\text{--}170 \times 30\text{--}40 \mu\text{m}$, regularly muriform; ascomata pseudostromatic, pseudostromata erumpent, with orange cover; Brazil. *Astrothelium carassense* Lücking, M.P. Nelsen & Marcelli

- N22b Ascospores $90\text{--}110 \times 22\text{--}28 \mu\text{m}$, with few longitudinal septa, submuriform; ascomata solitary to irregularly grouped, covered by thallus except ostiole; neotropical. *Astrothelium diplocarpum* Nyl.
- N23a Ascospores $100\text{--}130 \times 25\text{--}30 \mu\text{m}$, 8 per ascus; ascomata solitary, immersed to erumpent, irregularly bumpy with central red ostiole; neotropical. *Astrothelium purpurascens* (Müll. Arg.) Aptroot & Lücking
- N23b Ascospores $150\text{--}200 \times 40\text{--}50 \mu\text{m}$, 2 per ascus; Brazil. *Astrothelium isabellinum* Eschw.

Key O. *Astrothelium* species with ascospores muriform, ostiole lateral, secondary substances absent

- O1a Ascomata solitary, with separate ostioles. 2
- O1b Ascomata fused, with joint ostioles. 18
- O2a Hamathecium inspersed. 3
- O2b Hamathecium clear. 11
- O3a Ascospores $30\text{--}33 \times 9.5\text{--}10.5 \mu\text{m}$; Brazil. *Astrothelium quasimamillanum* Aptroot & C.O. Mendonça
- O3b Ascospores $70\text{--}230 \times 20\text{--}55 \mu\text{m}$. 4
- O4a Ascospores $70\text{--}100 \times 20\text{--}35 \mu\text{m}$; thallus lacking pseudocyphellae. 5
- O4b Ascospores $(85\text{--})100\text{--}230 \times 25\text{--}55 \mu\text{m}$, if less than $100 \mu\text{m}$ then thallus with pseudocyphellae. 7
- O5a Ascospores curved, $74\text{--}90 \times 25\text{--}34 \mu\text{m}$; ascomata dispersed, immersed to erumpent and covered by thallus except ostioles; Brazil. *Astrothelium curvatum* Aptroot & M. Cáceres
- O5b Ascospores straight. 6
- O6a Ascospores 8 per ascus; thallus smooth to uneven; ascomata dispersed to irregularly confluent, immersed to erumpent, laterally covered by thallus but otherwise exposed and black, surrounded by irregular whitish rim; pantropical. *Astrothelium santessonii* (Letr.-Gal.) Aptroot & Lücking
- O6b Ascospores 2 per ascus; thallus verrucose-papillose; ascomata dispersed to irregularly confluent, sessile, covered by thallus except the strongly protruding ostiole; Costa Rica. *Astrothelium minicecidiogenum* Aptroot & Sipman

- O7a Ascospores $250\text{--}350 \times 40\text{--}60 \mu\text{m}$, 1 per ascus; thallus irregularly bullate-bumpy; ascomata solitary, immersed, hardly visible; Panama. *Astrothelium megacrypticum* Lücking, M.P. Nelsen & N. Salazar
- O7b Ascospores $85\text{--}230 \times 25\text{--}60 \mu\text{m}$, (1–)2–8 per ascus; ascomata variable. 8
- O8a Ascospores 8 per ascus, $200\text{--}230 \times 30\text{--}40 \mu\text{m}$, about 6–7 times as long as broad; Brazil. *Astrothelium longisporum* Aptroot, J.R. Silva & M. Cáceres
- O8b Ascospores 1–2 per ascus. 9
- O9a Ascospores $170\text{--}340 \times 30\text{--}60 \mu\text{m}$, about 5–6 times as long as broad, 2 per ascus; neotropical. *Astrothelium puiggarii* (Müll. Arg.) Aptroot & Lücking
- O9b Ascospores $85\text{--}175 \times 25\text{--}55 \mu\text{m}$, about 2.5–4 times as long as broad. 10
- O10a Thallus with pseudocyphellae; ascomata laterally covered by thallus, with black ostiolar area surrounded by black rim; Bolivia. *Astrothelium pyrenuliforme* Flakus & Aptroot
- O10b Thallus lacking pseudocyphellae; ascomata solitary, exposed and pure black; Ecuador. *Astrothelium ecuadoriense* Aptroot
- O11a Ascospores 210–230 μm long, 1 per ascus; thallus irregularly verrucose; ascomata dispersed, immersed, hardly visible; São Tomé. *Astrothelium lugescens* (Nyl.) Aptroot & Lücking
- O11b Ascospores 70–200 μm long, (1–)2–8 per ascus; ascomata variable. 12
- O12a Ascospores $70\text{--}130 \times 20\text{--}36 \mu\text{m}$. 13
- O12b Ascospores $120\text{--}200 \times 30\text{--}50(\text{--}60) \mu\text{m}$. 15
- O13a Ascospores 2 per ascus; ascomata irregularly confluent to pseudostromatic, immersed to erumpent, covered by thallus except black ostiolar area surrounded by irregular whitish rim; India. *Astrothelium andamanicum* (Makhija & Patw.) Aptroot & Lücking
- O13b Ascospores 4–8 per ascus; ascomata dispersed. 14
- O14a Ascospores with distinctly thickened median septum, 4–8 per ascus; thallus smooth to uneven; ascomata erumpent; pantropical. *Astrothelium tenue* (Aptroot) Aptroot & Lücking
- O14b Ascospores lacking thickened median septum, 4 per ascus; thallus gall-like, coarsely bullate to folded; ascomata solitary, immersed, visible only by the dark ostiolar area; Guyana. *Astrothelium guianense* Aptroot
- O15a Ascospores I+ violet; thallus strongly bullate; ascomata immersed in thallus warts, completely covered by thallus; Bolivia. *Astrothelium bullatum* Flakus & Aptroot

- O15b Ascospores I–. 16
- O16a Ascospores 8 per ascus; eastern palaeotropical. *Astrothelium superbum* (Fr.) Aptroot & Lücking
- O16b Ascospores 2–4 per ascus. 17
- O17a Ascomata usually single in thalline warts, warts up to 1 mm diam., with broad, diffusely grey-black ostiolar area; ostioles always eccentric; neotropical. *Astrothelium campylocartilagineum* Aptroot & Lücking
- O17b Ascomata 1–5 in thalline warts, warts up to 2 mm diam., with narrow, brown-black ostiolar area surrounded by a whitish rim; ostioles pointing in various directions including apical ones; Nicaragua. *Astrothelium nicaraguense* Lücking, M.P. Nelsen & T. Orozco
- O18a Ascospores small to medium-sized (30–80 × 16–26 µm). 19
- O18b Ascospores large (125–230 × 28–80 µm). 23
- O19a Hamathecium inspersed; ascospores 50–65 µm long; groups of fused ascomata irregularly confluent to pseudostromatic, immersed to erumpent, covered by thallus except black ostiolar area surrounded by irregular whitish rim; Brazil. *Astrothelium testudineum* Aptroot & M. Cáceres
- O19b Hamathecium clear; ascospores 30–52 or 70–80 µm long. 20
- O20a Ascospores medium-sized (70–80 × 20–25 µm); thallus uneven-verrucose; groups of fused ascomata pseudostromatic, pseudostromata prominent, covered by thallus except for dark, papilliform ostiole; Brazil. *Astrothelium mediocrassum* Aptroot
- O20b Ascospores small (30–52 × 16–22 µm); thallus and ascomata variable. 21
- O21a Groups of fused ascomata hemispherical; Brazil. *Astrothelium citrisporum* Aptroot, Oliveira-Junior & M. Cáceres
- O21b Groups of fused ascomata immersed to erumpent. 22
- O22a Groups of fused ascomata pseudostromatic, pseudostromata immersed, with white cover, forming irregular to reticulate lines, contrasting with thallus; neotropical. *Astrothelium cinereum* (Müll. Arg.) Aptroot & Lücking
- O22b Groups of fused ascomata dispersed, immersed-erumpent, hardly visible; neotropical. *Astrothelium leioplacum* (Müll. Arg.) Aptroot & Lücking
- O23a Ascospores 215–230 µm long, 1 per ascus; groups of fused ascomata dispersed, erumpent, completely covered by thallus; on rock; Brazil. *Astrothelium saxicola* (Malme) Aptroot & Lücking
- O23b Ascospores 125–200 µm long, 8 per ascus. 24

- O24a Ascospores 2/ascus, with median thickening; hamathecium only inspersed above; Brazil. *Astrothelium dimidioinspersum* Aptroot & M. Cáceres
- O24b Ascospores 8/ascus, with or without median thickening. 25
- O25a Ascospores with median thickening; groups of fused ascomata dispersed, immersed, hardly visible; Brazil. *Astrothelium flavomurisorum* Aptroot & M. Cáceres
- O25b Ascospores lacking thickening; groups of fused ascomata dispersed to irregularly confluent, covered by thallus except whitish ostiolar area; neotropical. *Astrothelium cecidiogenum* (Aptroot & Lücking) Aptroot & Lücking

Key P. *Bathelium* species

- P1a Ascospores transversely septate. 2
- P1b Ascospores muriform. 7
- P2a Ascospores 13–17-septate, 65–105 µm long; ostiolar area UV+ yellow, with lichexanthone; Bolivia. *Bathelium flavostiolum* Flakus & Aptroot
- P2b Ascospores 3–9-septate, 18–40 µm long; UV-. 3
- P3a Ascospores 3(–5)-septate, 18–25(–30) µm long. 4
- P3b Ascospores 5–9-septate, 25–40 µm long. 6
- P4a Internal pseudostroma pigment yellow, K+ persistently yellow; ascomata distinctly pseudostromatic; USA. *Bathelium carolinianum* (Tuck.) R.C. Harris
- P4b Internal pseudostroma pigment yellow-orange to reddish brown, K+ orange-red or purple-violet; ascomata variable. 5
- P5a Internal pseudostroma pigment yellow-orange, K+ purple-violet; ascospores 18–21 × 6–7 µm long; ascomata pseudostromatic; South Africa. *Bathelium austroafricanum* (Zahlbr.) Aptroot & Lücking
- P5b Internal pseudostroma pigment reddish brown, K+ orange-red; ascospores 20–30 × 6–7 µm long; ascomata solitary to irregularly confluent; eastern palaeotropical. *Bathelium porinosporum* Lücking, M.P. Nelsen & Gueidan
- P6a Ostioles usually white-pruinose; ascospores 5–7-septate, 25–32 µm long; eastern palaeotropical. *Bathelium albidoporum* (Makhija & Patw.) R.C. Harris
- P6b Ostioles non-pruinose, black; ascospores 5–9-septate, 30–40 µm long; palaeotropical. *Bathelium nigroporum* (Makhija & Patw.) Aptroot & Lücking

- P7a Thallus and pseudostromata UV+ yellow, with lichexanthone. 8
 P7b Thallus and pseudostromata UV- or UV+ red when pigment is present, lichexanthone absent. 9
- P8a Hamathecium inspersed; Brazil. *Bathelium pruinolucens* Aptroot & Lücking
 P8b Hamathecium not inspersed; Bolivia and Brazil. *Bathelium pruinosum* Flakus, Kukwa & Aptroot
- P9a Internal pigment of pseudostromata dark red, K+ green. 10
 P9b Internal pigment of pseudostromata yellow-orange, K+ purple. 11
- P10a Pseudostromata erumpent, covered with orange pigment and with dark ostioles surrounded by paler rim; Bolivia and Brazil. *Bathelium mirabile* Flakus, Kukwa & Aptroot
 P10b Pseudostromata prominent to sessile, dark brown and with strongly contrasting, whitish ostiolar area; Bolivia. *Bathelium boliviense* Flakus & Aptroot
- P11a Ascospores 40–80 µm long. 12
 P11b Ascospores 100–200 µm long. 15
- P12a Hamathecium fully inspersed; Bolivia. *Bathelium inspersionastoides* Flakus & Aptroot
 P12b Hamathecium clear except for ostiolar area. 13
- P13a Ascospores (50–)60–80 µm long. *Bathelium mastoideum* Afzel. ex Ach.
 P13b Ascospores 40–55 µm long. 14
- P14a Thallus smooth to uneven; pantropical. *Bathelium madreporiforme* (Eschw.) Trevis.
 P14b Thallus verrucose-bullate; pantropical. *Bathelium tuberculosum* (Makhija & Patw.) R.C. Harris
- P15a Ascospores 100–120 µm long; pantropical. *Bathelium lineare* (C.W. Dodge) R.C. Harris
 P15b Ascospores 130–200 µm long; Sierra Leone. *Bathelium sphaericum* (C.W. Dodge) R.C. Harris

Key Q; pantropical. *Bogoriella* species

- Q1a Ascospores becoming distinctly muriform, with thin eusepta and rectangular lumina.
 2

- Q1b Ascospores 1-septate to rarely 3-septate or submuriform, with thin eusepta or with irregularly thickened endospore and distosepta. 4
- Q2a Ascospores becoming thick-walled and with thickened median septum, 28–45 μm long; Bolivia. *Bogoriella chiquitana* (Flakus, Kukwa & Aptroot) Aptroot & Lücking
- Q2b Ascospores remaining thin-walled, 20–35 μm long. 3
- Q3a Ascospores 20–30 \times 12–16 μm , about 1.5–2 times as long as broad; pantropical. *Bogoriella decipiens* (Müll. Arg.) Aptroot & Lücking
- Q3b Ascospores 20–35 \times 7–10 μm , about 3–3.5 times as long as broad; China. *Bogoriella pandanicola* (S.N. Zhang & K.D. Hyde) S.N. Zhang, Aptroot & Lücking
- Q4a Ascospores with irregularly thickened endospore and distosepta, 1-septate or rarely 3-septate to submuriform; lumina irregular to halter-shaped. 5
- Q4b Ascospores with thin eusepta, 1-septate or rarely 3-septate; lumina rectangular. 8
- Q5a Ascospores (1–)3-septate to appearing submuriform, 27–45 \times 10–17 μm . 6
- Q5b Ascospores 1-septate but cells with irregular endospore invaginations, 15–29 \times 6–14 μm . 7
- Q6a Ascospores mostly distinctly 3-septate, occasionally 1-septate or with additional longitudinal septa; pantropical. *Bogoriella queenslandica* (Müll. Arg.) Aptroot & Lücking
- Q6b Ascospores with one distinct, rather dark central septum and the two cells with irregular endospore invaginations giving them the appearance of a (sub-)muriform division; Brazil. *Bogoriella complexoluminata* Aptroot & Lücking
- Q7a Ostiole with red pigment; ascospores 23–29 \times 10–14 μm , secondary endospore invaginations developing late; Hong Kong. *Bogoriella isthmospora* (Aptroot) Lücking, R. Miranda & Aptroot
- Q7b Ostiole lacking pigment; ascospores 15–18 \times 6–8 μm , secondary endospore invaginations developing soon; Caribbean. *Bogoriella rubrostoma* (Aptroot) Aptroot & Lücking
- Q8a Ostiole apical. 9
- Q8b Ostiole lateral. 18
- Q9a Hamathecium inspersed. 10
- Q9b Hamathecium clear. 12

-
- Q10a Ascospores $24\text{--}27 \times 8\text{--}11 \mu\text{m}$; neotropical. *Bogoriella oleosa* (Aptroot) Aptroot & Schumm
- Q10b Ascospores $30\text{--}45 \times 10\text{--}15 \mu\text{m}$. 11
- Q11a Ascomata 0.5–0.7 mm diam; Venezuela. *Bogoriella xanthonica* (Komposch, Aptroot & Hafellner) Aptroot & Lücking
- Q11b Ascomata 0.7–1.1 mm diam; Venezuela. *Bogoriella macrocarpa* (Komposch, Aptroot & Hafellner) Aptroot & Lücking
- Q12a Ascospores becoming 3-septate; pantropical. *Bogoriella obovata* (Stirt.) Aptroot & Lücking
- Q12b Ascospores remaining 1-septate. 13
- Q13a Ascospores 28–40 μm long; Brazil. *Bogoriella megaspora* (Aptroot & M. Cáceres) Aptroot & Lücking
- Q13b Ascospores 17–27 μm long. 14
- Q14a Black prothallus line present. 15
- Q14b Black prothallus line absent. 16
- Q15a Asci 10–20 μm broad; eastern palaeotropical. *Bogoriella conothelena* (Nyl.) Aptroot & Lücking
- Q15b Asci 20–30 μm broad; pantropical. *Bogoriella thelena* (Ach.) Aptroot & Lücking
- Q16a Ascomata 0.8–1 mm diam., basally broadly expanded; pantropical. *Bogoriella confuens* (Müll. Arg.) Aptroot & Lücking
- Q16b Ascomata 0.2–0.4 mm diam., basally barely or not expanded. 17
- Q17a Ascomata conical; Brazil. *Bogoriella pachythea* (Sacc. & P. Syd.) Aptroot & Lücking
- Q17b Ascomata fattened; pantropical. *Bogoriella modesta* (Müll. Arg.) Aptroot & Lücking
- Q18a Thallus UV-negative; Puerto Rico. *Bogoriella triangularis* (Aptroot) Aptroot & Lücking
- Q18b Thallus UV+ yellow; Brazil. *Bogoriella xantholateralis* Aptroot

Key R. *Constrictolumina* species

- R1a Ascospores $17\text{--}23 \times 5\text{--}8 \mu\text{m}$, 1–3-septate. 2
- R1b Ascospores (20–)22–46 \times 7–24 μm , 1-septate. 3

- R2a Ascomata aggregate; neotropical. *Constrictolumina esenbeckiana* (Fée) Lücking, M.P. Nelsen & Aptroot
- R2b Ascomata solitary; pa; Dominica; neotropical ntropical. *Constrictolumina cinchonae* (Ach.) Lücking, M.P. Nelsen & Aptroot
- R3a Ascospores $40\text{--}46 \times 16\text{--}24 \mu\text{m}$. *Constrictolumina porospora* (Vain.) Lücking, M.P. Nelsen & Aptroot
- R3b Ascospores $20\text{--}30 \times 7\text{--}16 \mu\text{m}$. 4
- R4a Ascomata aggregate; ascospores 3-septate; neotropical. *Constrictolumina leucostoma* (Müll. Arg.) Lücking, M.P. Nelsen & Aptroot
- R4b Ascomata solitary; ascospores 1(–3)-septate; pantropical. *Constrictolumina planorbis* (Ach.) Lücking, M.P. Nelsen & Aptroot

Key S. *Dictyomeridium* species

- S1a Thallus UV+ yellow. 2
- S1b Thallus UV–. 5
- S2a Ascomata deeply immersed in the bark, ascospores 2 per ascus, $45\text{--}70 \times 17\text{--}25 \mu\text{m}$; Brazil. *Dictyomeridium immersum* (Aptroot, A.A. Menezes & M. Cáceres) Aptroot, M.P. Nelsen & Lücking
- S2b Ascomata erumpent, ascospores 8 per ascus. 3
- S3a Ascospores $25\text{--}35 \times 12\text{--}17 \mu\text{m}$; Bolivia. *Dictyomeridium lueckingii* Flakus & Aptroot
- S3b Ascospores over $35 \mu\text{m}$ long. 4
- S4a Ascospores $35\text{--}54 \times 12\text{--}19 \mu\text{m}$; pantropical. *Dictyomeridium proponens* (Nyl.) Aptroot, M.P. Nelsen & Lücking
- S4b Ascospores $55\text{--}75 \times 19\text{--}21 \mu\text{m}$; Brazil. *Dictyomeridium parapropionens* (Aptroot, M. Cáceres & E.L. Lima) Aptroot, M.P. Nelsen & Lücking
- S5a Ascospores $55\text{--}80 \times 17\text{--}25 \mu\text{m}$; eastern palaeotropical. *Dictyomeridium campylothelioides* (Aptroot & Sipman) Aptroot, M.P. Nelsen & Lücking
- S5b Ascospores under $55 \mu\text{m}$ long. 6
- S6a Ostiole with red, KOH+ green pigment inside, ascospores $33\text{--}55 \times 11\text{--}15 \mu\text{m}$; Brazil. *Dictyomeridium isohypocrellinum* (Xavier-Leite, M. Cáceres & Aptroot) Aptroot, M.P. Nelsen & Lücking

S6b Ostiole without red pigment, ascospores 33–55 × 12–21 µm; pantropical.
Dictyomeridium amylosporium (Vain.) Aptroot, M.P. Nelsen & Lücking

Key T. *Macroconstrictolumina* species

T1a Ascospores 18–30 × 7–10(–12) µm; pantropical. *Macroconstrictolumina lyrata*
(R.C. Harris) Lücking, R. Miranda & Aptroot

T1b Ascospores 30–48 × 10–16(–24) µm. 2

T2a Ostiole lateral; ascospores 1-septate; Brazil. *Macroconstrictolumina megalateralis*
Aptroot

T2b Ostiole apical; ascospores 1(–3)-septate. 3

T3a Ascospores 27–37 × 10–12 µm, often 3-septate; pantropical. *Macroconstrictolumina*
majuscula (Nyl.) Lücking, R. Miranda & Aptroot

T3b Ascospores 40–50 × 15–20 µm, mostly 1-septate; pantropical.

Macroconstrictolumina malaccitula (Nyl.) Lücking, R. Miranda & Aptroot

Key U. *Marcelaria* species

U1a Ascomata (and often thallus) bright red, UV–; hamathecium clear; ascospores 110–
170 µm long, amyloid; Neotropics and African Palaeotropics. *Marcelaria*
purpurina (Nyl.) Aptroot, Nelsen & Parmen

U1b Ascomata (and often thallus) (orange-)yellow, UV+ yellow; hamathecium densely
inspersed; ascospores 50–80 µm long, non-amyloid (rarely dextrinoid); eastern
Palaeotropics. 2

U2a Thallus yellow-pruinose; ascospores 8 per ascus; ostiolar region narrow. *Marcelaria*
benguelensis (Müll. Arg.) Aptroot, Nelsen & Parmen

U2b Thallus non-pruinose; ascospores 2 per ascus; ostiolar region broad, flat. *Marcelaria*
cumingii (Mont.) Aptroot, Nelsen & Parmen

Key V. *Nigrovothelium* species

V1a Hamathecium inspersed; pantropical. *Nigrovothelium inspersotropicum* Aptroot &
Diederich

V1b Hamathecium clear. 2

- V2a Thallus smooth to uneven; pantropical. *Nigrovothelium tropicum* (Ach.) Lücking, M.P. Nelsen & Aptroot
V2b Thallus verrucose-bullate; India. *Nigrovothelium bullatum* Lücking, Upreti & Lumbsch

Key W. *Polymeridium* species

- W1a Ascospores transversely septate. 2
W1b Ascospores submuriform to muriform. 49
- W2a Ascospores regularly 3-septate. 3
W2b Ascospores irregularly (3–)4–17-septate. 28
- W3a Thallus UV+ yellow, with lichexanthone. 4
W3b Thallus UV–, without substances or rarely with medullary pigments or pruina on ascomata. 14
- W4a Hamathecium interspersed. 5
W4b Hamathecium clear. 7
- W5a Ascospores 24–33 × 6–10 µm; pantropical. *Polymeridium subvirescens* (Nyl.) R.C. Harris
W5b Ascospores 17–23 × 5–8 µm. 6
- W6a Ascospores 17–20 × 5–7 µm; pantropical. *Polymeridium pyrenuloides* (Müll. Arg.) Aptroot
W6b Ascospores 20–23 × 6–8 µm; pantropical. *Polymeridium albidovarians* Aptroot
- W7a Ostioles eccentric, separate or fused; ascospores 24–33 µm long. 8
W7b Ostioles apical; ascospores variable. 9
- W8a Ostioles fused; Venezuela. *Polymeridium pyrenastroides* R.C. Harris ex Aptroot
W8b Ostioles separate; neotropical. *Polymeridium stramineoatrum* (Vain.) Aptroot
- W9a Ascospores IKI+ violet, 16–23 × 5–7 µm; Brazil. *Polymeridium amyloideum* R.C. Harris
W9b Ascospores IKI–. 10
- W10a Ascospores 13–20 × 4–6 µm. 11
W10b Ascospores 19–33 × 6–11 µm. 12

- W11a Ascospores 13–17 μm long; pantropical. *Polymeridium albopruinosum* (Makhija & Patw.) Aptroot
- W11b Ascospores 16–20 μm long; pantropical. *Polymeridium bengoanum* (Vain.) Aptroot
- W12a Ascomata mostly covered by thallus, with black ostioles bordered by white ring; ascospores relatively broad, 17–25 \times 7–11 μm , c. 2.5 times as long as broad; Australia. *Polymeridium xanthoreagens* Aptroot
- W12b Ascomata pure black; ascospores narrower, 20–33 \times 6–10 μm , about 3 times as long. 13
- W13a Ascospores 20–23 \times 6–8 μm ; Brazil. *Polymeridium albidoreagens* Aptroot, A.A. Menezes & M. Cáceres
- W13b Ascospores 25–33 \times 7–10 μm ; pantropical. *Polymeridium catapastum* (Nyl.) R.C. Harris
- W14a Hamathecium inspersed; ostioles apical. 15
- W14b Hamathecium clear; ostioles variable. 20
- W15a Ascospores 24–34 μm long; ascomata laterally covered by thallus. 16
- W15b Ascospores 12–24 μm long; ascomata variable. 17
- W16a Ascospores with rough wall, 26–34 μm long; blackish ostiolar area broad; eastern palaeotropical. *Polymeridium glaucoatrum* (Vain.) R.C. Harris
- W16b Ascospores smooth, 24–30 μm long; blackish ostiolar area narrow; eastern palaeotropical. *Polymeridium suffusum* (Knight) Aptroot
- W17a Ascospores 12–20 \times 3–6 μm . 18
- W17b Ascospores 20–24 \times 5–10 μm . 19
- W18a Ascospores 12–16 μm long; neotropical. *Polymeridium contendens* (Nyl.) R.C. Harris
- W18b Ascospores 16–20 μm long; pantropical. *Polymeridium inspersionum* Aptroot
- W19a Ascomata with ostioles bordered by white ring; ascospores relatively broad, 20–24 \times 7–10 μm , c. 2.5–3 times as long as broad; pantropical. *Polymeridium tribulationis* Aptroot
- W19b Ascomata pure black; ascospores narrower, 20–23 \times 5–7 μm , c. 3–4 times as long as broad; pantropical. *Polymeridium siamense* (Vain.) Aptroot
- W20a Ascomata with internal or external, yellow-orange or red pigment. 21

- W20b Ascomata without pigments. 22
- W21a Ascomata with internal, yellow-orange, K+ purple pigment; ascospores $30\text{--}34 \times 9\text{--}10 \mu\text{m}$; Brazil. *Polymeridium endocrocinum* R.C. Harris
- W21b Ascomata with external, red, K+ red pruina; ascospores $17\text{--}19 \times 4\text{--}5 \mu\text{m}$; Puerto Rico. *Polymeridium rhodopruinosum* Aptroot & Mercado Diaz
- W22a Ostioles lateral. 23
- W22b Ostioles apical. 24
- W23a Ascomata grouped with fused ostioles; ascospores $26\text{--}30 \times 8\text{--}10 \mu\text{m}$; Brazil. *Polymeridium biloculare* R.C. Harris
- W23b Ascomata solitary with separate ostioles; ascospores $30\text{--}35 \times 12\text{--}13 \mu\text{m}$; neotropical. *Polymeridium simulans* R.C. Harris
- W24a Ascospores $24\text{--}33 \mu\text{m}$ long; eastern palaeotropical. *Polymeridium catapastoides* Aptroot
- W24b Ascospores $11\text{--}27 \mu\text{m}$ long. 25
- W25a Ascospores $11\text{--}16 \times 3\text{--}6 \mu\text{m}$; pantropical. *Polymeridium microsporum* (Makhija & Patw.) Aptroot
- W25b Ascospores longer. 26
- W26a Ascospores $(15\text{--})16\text{--}20 \times 5.0\text{--}7.5 \mu\text{m}$; pantropical. *Polymeridium subcinereum* (Nyl.) R.C. Harris
- W26b Ascospores $19\text{--}27 \times 6\text{--}11 \mu\text{m}$. 27
- W27a Ostiole flat and with white ring; ascospores $19\text{--}27 \times 6.5\text{--}11.0 \mu\text{m}$; eastern palaeotropical.
Polymeridium sulphurescens (Müll. Arg.) R.C. Harris
- W27b Ostiole not flat, black; ascospores $(18\text{--})20\text{--}23 \times 6\text{--}8(-9) \mu\text{m}$; pantropical.
Polymeridium albidum (Müll. Arg.) R.C. Harris
- W28a Thallus UV+ yellow, with lichexanthone. 29
- W28b Thallus UV-, without substances. 38
- W29a Hamathecium inspersed. 30
- W29b Hamathecium clear. 33
- W30a Hamathecium yellow. 31
- W30b Hamathecium hyaline. 32

- W31a Ascospores 5–7-septate, 32–37 × 10–13 µm; Brazil. *Polymeridium endoflavens* Aptroot, D.S. Andrade & M. Cáceres
- W31b Ascospores 8–12-septate, 56–70 × 12–14 µm; Brazil. *Polymeridium longiflavens* Aptroot, Mendonça & M. Cáceres
- W32a Ascospores 4–7-septate, 18–28 × 5–7 µm; pantropical. *Polymeridium multiseptatum* Aptroot, A.A. Menezes & M. Cáceres
- W32b Ascospores 8–12-septate, 40–50 × 9–11 µm; neotropical. *Polymeridium chioneum* (Mont.) R.C.Harris
- W33a Ascomata with lateral ostioles. 34
- W33b Ascomata with apical ostioles. 36
- W34a Ascomata solitary, ascospores 11–13-septate, 52–88 × 16–19 µm; Panama. *Polymeridium xanthopleurothecium* Aptroot & Etayo
- W34b Ascomata grouped with lateral, fused ostioles. 35
- W35a Ascospores 28–35 µm long; neotropical. *Polymeridium dithecium* R.C. Harris
- W35b Ascospores 30–50 µm long; Bolivia. *Polymeridium xanthoexcentricum* Flakus & Aptroot
- W36a Ascospores (4–)5(–7)-septate, 19–26 µm long; neotropical. *Polymeridium multiforme* Aptroot
- W36b Ascospores 7–13-septate, 28–39 µm long. 37
- W37a Thallus corticate; Brazil. *Polymeridium corticatum* A.A. Menezes, M. Cáceres & Aptroot
- W37b Thallus ecorticate; neotropical. *Polymeridium alboflavescens* Aptroot
- W38a Hamathecium inspersed. 39
- W38b Hamathecium clear. 45
- W39a Hamathecium yellow. 40
- W39b Hamathecium hyaline. 41
- W40a Ascospores 8–12-septate, 40–55 × 9–11 µm; ostioles lateral; ; neotropical. *Polymeridium flavoethecium* R.C. Harris
- W40b Ascospores 6–8-septate, 35–37 × 7.5–8 µm; ostioles apical; Sri Lanka. *Polymeridium fernandoi* Aptroot & Weerakoon
- W41a Ostioles lateral; ascospores 3–6-septate, 22–26 × 7–9 µm; India. *Polymeridium refertum* (Stirt.) Aptroot

- W41b Ostioles apical; ascospores variable. 42
- W42a Ascospores 4–7-septate. 43
W42b Ascospores 7–12-septate. 44
- W43a Ascospores with rough wall, 7-septate, $30\text{--}33 \times 9\text{--}10 \mu\text{m}$; Venezuela.
Polymeridium neuwirthii Aptroot
W43b Ascospores smooth, (4–)5(–7)-septate, $18\text{--}28 \times 4\text{--}7 \mu\text{m}$; pantropical.
Polymeridium quinquesepatum (Nyl.) R.C. Harris
- W44a Ascospores 7–11-septate, $25\text{--}36 \times 5\text{--}9 \mu\text{m}$; pantropical. *Polymeridium pleiomerellum* (Müll. Arg.) R.C. Harris
W44b Ascospores 8–12-septate, $40\text{--}50 \times 9\text{--}11 \mu\text{m}$; neotropical. *Polymeridium costaricense* Aptroot
- W45a Ostioles lateral. 46
W45b Ostioles apical. 47
- W46a Ascospores 7–9-septate, $40\text{--}50 \times 8\text{--}9 \mu\text{m}$; neotropical. *Polymeridium neblinae* R.C. Harris
W46b Ascospores 9–13-septate, $(35\text{--})50\text{--}75 \times 12\text{--}16 \mu\text{m}$; pantropical. *Polymeridium pleurothecium* R.C. Harris
- W47a Ascospores 13–17-septate, $85\text{--}150 \times 20\text{--}30 \mu\text{m}$; Argentina. *Polymeridium bambusicola* Aptroot & L.I. Ferraro
W47b Ascospores 4–11-septate, $21\text{--}39 \times 5\text{--}9 \mu\text{m}$. 48
- W48a Ascospores (4–)5(–7)-septate, $21\text{--}30 \times 5\text{--}9 \mu\text{m}$; pantropical. *Polymeridium jordanii* (C.W. Dodge) Aptroot
W48b Ascospores 7–11-septate, $(24\text{--})28\text{--}39 \times 6\text{--}9 \mu\text{m}$; pantropical. *Polymeridium albocinereum* (Kremp.) R.C. Harris
- W49a Hamathecium inspersed, ascospores $18\text{--}20 \times 6.0\text{--}7.5 \mu\text{m}$; Philippines.
Polymeridium submuriforme Aptroot
W49b Hamathecium clear, ascospores $25\text{--}45 \times 9\text{--}13 \mu\text{m}$. 50
- W50a Ascospores $30\text{--}45 \times 9\text{--}11 \mu\text{m}$, submuriform, 3–4 times as long as broad; neotropical. *Polymeridium cinereonigricans* (Vain.) R.C. Harris
W50b Ascospores $25\text{--}29 \times 11\text{--}13 \mu\text{m}$, muriform, 2.0–2.5 times as long as broad; Brazil.
Polymeridium julelloides E.L. Lima, M. Cáceres & Aptroot

Key X. *Pseudobogoriella* species

X1a Ostiole lateral; Australia. *Pseudobogoriella lateralis* (Sipman) Lücking, R. Miranda & Aptroot

X1b Ostiole apical 2

X2a Ascospores 24–28 µm long; pantropical. *Pseudobogoriella hemisphaerica* (Müll. Arg.) Lücking, R. Miranda & Aptroot

X2b Ascospores 12–22 µm long. 3

X3a Thallus foliicolous, subcuticular, distinctly lobed; New Zealand. *Pseudobogoriella striguloides* (Sérus. & Aptroot) Lücking, R. Miranda & Aptroot

X3b Thallus corticolous, not lobed. 4

X4a Excipulum with K⁺ red, dissolving pigment; pantropical. *Pseudobogoriella leuckertii* (D. Hawksw. & J.C. David) Lücking, R. Miranda & Aptroot

X4b Excipulum lacking pigment. 5

X5a Black prothallus line with numerous pycnidia present; pantropical. *Pseudobogoriella punctata* (Aptroot) Lücking, R. Miranda & Aptroot

X5b Black prothallus line lacking pycnidia or absent. 6

X6a Thallus besides ascomata with numerous black dots representing pycnidia. 7

X6b Thallus lacking pycnidia. 8

X7a Ascomata aggregate; ascospores 15–20 × 6–9 µm; neotropical. *Pseudobogoriella socialis* (Zahlbr.) Lücking, R. Miranda & Aptroot

X7b Ascomata solitary; ascospores 11–17 × 5–7 µm; pantropical. *Pseudobogoriella fumosula* (Zahlbr.) Lücking, R. Miranda & Aptroot

X8a Involucrellum broadly expanding basally to over 0.5 mm diam; pantropical.

Pseudobogoriella alata (Groenh. ex Aptroot) Lücking, R. Miranda & Aptroot

X8b Ascomata including basally slightly expanding involucrellum 0.2–0.5 mm diam. 9

X9a Ascospores 12–16 µm long. 10

X9b Ascospores 16–22 µm long. 12

X10a Ascomata 03–05 mm diam; pantropical. *Pseudobogoriella subfallens* (Müll. Arg.) Lücking, R. Miranda & Aptroot

X10b Ascomata 02–03 mm diam. 11

- X11a Ascospores 4–55 μm broad, 2.5–3 times as long as broad; involucrellum basally not expanding; pantropical. *Pseudobogoriella exigua* (Müll. Arg.) Lücking, R. Miranda & Aptroot
- X11b Ascospores 5.5–7 μm broad, 2–2.5 times as long as broad; involucrellum basally expanding; China. *Pseudobogoriella minutula* (Zahlbr.) Lücking, R. Miranda & Aptroot
- X12a Black prothallus line present. 13
- X12b Black prothallus line absent. 14
- X13a Ascomata conical, basally not expanded; neotropical. *Pseudobogoriella captiosa* (Müll. Arg.) Lücking, R. Miranda & Aptroot
- X13b Ascomata fattened, basally expanded; pantropical. *Pseudobogoriella miculiformis* (Müll. Arg.) Lücking, R. Miranda & Aptroot
- X14a Ascomata basally broadly expanded, fringe 50–100 μm Broad; pantropical. *Pseudobogoriella annonacea* (Müll. Arg.) Lücking, R. Miranda & Aptroot
- X14b Ascomata basally narrowly expanded, fringe 20–50 μm Broad; pantropical. *Pseudobogoriella nonensis* (Stirt.) Lücking, R. Miranda & Aptroot

Key Y. *Pseudopyrenula* species

- Y1a Thallus UV+ yellow, with lichexanthone. 2
- Y1b Thallus UV-. 6
- Y2a Hamathecium clear; ascospores 30–45 \times 8–11 μm ; Venezuela. *Pseudopyrenula cryptotheca* Komposch, Aptroot & Hafellner
- Y2b Hamathecium interspersed with hyaline or yellow oil droplets; ascospores under 37 μm long. 3
- Y3a Hamathecium interspersed with yellow oil droplets. 4
- Y3b Hamathecium interspersed with hyaline oil droplets. 5
- Y4a Ascospores 21–28 \times 6–9 μm ; neotropical. *Pseudopyrenula flavoreagens* Aptroot & M. Cáceres
- Y4b Ascospores 26–36 \times 7–10 μm ; Bolivia. *Pseudopyrenula flavosuperans* Flakus & Aptroot
- Y5a Ascospores 21–25 \times 6–9 μm ; neotropical. *Pseudopyrenula guianensis* Aptroot

- Y5b Ascospores $30\text{--}35 \times 9\text{--}11 \mu\text{m}$; neotropical. *Pseudopyrenula superans* Müll. Arg.
- Y6a Ascospores 5-septate, $16\text{--}21 \times 6\text{--}7 \mu\text{m}$; Venezuela. *Pseudopyrenula hexamera*
Aptroot
- Y6b Ascospores 3-septate. 7
- Y7a Thallus corticate; ascospores $21\text{--}25 \times 6\text{--}9 \mu\text{m}$; Costa Rica. *Pseudopyrenula thallina*
Lücking & Aptroot
- Y7b Thallus not corticate. 8
- Y8a Hamathecium interspersed with hyaline or yellow oil droplets. 9
- Y8b Hamathecium clear (but ascospores may contain yellow oil). 19
- Y9a Hamathecium interspersed with yellow oil droplets. 10
- Y9b Hamathecium interspersed with hyaline oil droplets (but ascospores may contain
yellow oil). 14
- Y10a Ascospores $37\text{--}44 \times 10\text{--}12 \mu\text{m}$; neotropical. *Pseudopyrenula endoxantha* (Vain.)
Zahlbr.
- Y10b Ascospores under $35 \times 10 \mu\text{m}$. 11
- Y11a Ascospores $25\text{--}30 \times 7.5\text{--}8.5 \mu\text{m}$; pantropical. *Pseudopyrenula media* Aptroot &
Diederich
- Y11b Ascospores $13\text{--}25 \times 5.5\text{--}9 \mu\text{m}$. 12
- Y12a Ascospores $13\text{--}17 \times 5.5\text{--}6.5 \mu\text{m}$; Brazil. *Pseudopyrenula miniflavida* Aptroot &
A.D. Nunes
- Y12b Ascospores $18\text{--}25 \times 6\text{--}9 \mu\text{m}$. 13
- Y13a Ascospores $21\text{--}25 \times 6\text{--}9 \mu\text{m}$; hamathecium K+ deep yellow to reddish; pantropical.
Pseudopyrenula subgregaria Müll. Arg.
- Y13b Ascospores $18\text{--}22 \times 6\text{--}8 \mu\text{m}$; hamathecium K+ weakly purplish; Thailand.
Pseudopyrenula endoxanthoides Vain.
- Y14a Ascoma wall with yellow, K+ red pigment; ascospores $21\text{--}32 \times 7\text{--}10 \mu\text{m}$; Brazil.
Pseudopyrenula cerei Vain.
- Y14b Ascoma wall without yellow pigment. 15
- Y15a Thallus saxicolous; ascospores $35\text{--}45 \times 11\text{--}13 \mu\text{m}$; Brazil. *Pseudopyrenula*
saxicola Malme
- Y15b Thallus corticolous; ascospores under $38 \mu\text{m}$ long. 16

- Y16a Ascospores $33\text{--}37 \times 10\text{--}12 \mu\text{m}$; neotropical. *Pseudopyrenula dubia* Vain.
Y16b Ascospores under $33 \mu\text{m}$ long. 17
- Y17a Ascospores $(24\text{--})26\text{--}32 \times 7\text{--}10 \mu\text{m}$; pantropical. *Pseudopyrenula diluta* (Fée) Müll. Arg.
Y17b Ascospores $21\text{--}25 \times 6\text{--}9 \mu\text{m}$. 18
- Y18a Ostioles eccentric; Cuba. *Pseudopyrenula cubana* (Müll. Arg.) Aptroot & Lücking
Y18b Ostioles apical; pantropical. *Pseudopyrenula subnudata* Müll. Arg.
- Y19a Thallus without algae, brownish, of bark colour. 20
Y19b Thallus with algae, whitish. 22
- Y20a Ascospores $45\text{--}55 \times 12\text{--}15 \mu\text{m}$; USA (Florida). *Pseudopyrenula staphyleae* (Petr.) Aptroot
Y20b Ascospores under $40 \mu\text{m}$ long. 21
- Y21a Ascospores $25\text{--}29 \times 7\text{--}9 \mu\text{m}$; Papua New Guinea. *Pseudopyrenula papuana* Aptroot
Y21b Ascospores $30\text{--}39 \times 6\text{--}8 \mu\text{m}$; Papua New Guinea. *Pseudopyrenula serusiauxii* Aptroot
- Y22a Ascospores $12\text{--}18 \times 4\text{--}6 \mu\text{m}$; Tasmania. *Pseudopyrenula kantvilasii* (P.M. McCarthy) Aptroot, comb. nov. Mycobank no 841169; basionym: *Lithothelium kantvilasii* P.M. McCarthy, Telopea 18: 168, 2015. The description and illustration show several characters that are characteristic of the genus *Pseudopyrenula*, especially the presence of yellow oil in the ascospores, which is unknown from any other genus, but also the diamond-shaped lumina and the branched hamathecium filaments.
Y22b Ascospores $26\text{--}35 \times 7\text{--}11 \mu\text{m}$. 23
- Y23a Ascospores $26\text{--}32 \times 7\text{--}10 \mu\text{m}$; neotropical. *Pseudopyrenula americana* Aptroot
Y23b Ascospores $30\text{--}35 \times 9\text{--}11 \mu\text{m}$; Brazil. *Pseudopyrenula gelatinosa* Aptroot

Key *Z. Trypethelium* species

- Z1a Thallus, pseudostroma and/or ostioles UV+ yellow, with lichexanthone 2
Z1b Thallus, pseudostroma and ostioles UV-. 9
- Z2a Thallus UV-, only pseudostroma and/or ostioles UV+ yellow 3
Z2b Thallus and sometimes also pseudostroma UV+ yellow 4

- Z3a Pseudostroma UV+ yellow outside, with yellow pigment inside. *Trypethelium aureornatum* Aptroot
- Z3b Only ostioles UV+ yellow; Brazil. *Trypethelium xanthostiolornatum* Aptroot
- Z4a Hamathecium not inspersed. 5
- Z4b Hamathecium inspersed. 8
- Z5a Pseudostromata externally with orange pigment; Bolivia. *Trypethelium xanthoplatystomum* Flakus & Aptroot
- Z5b Pseudostromata internally with yellow pigment. 6
- Z6a Thallus and pseudostroma UV+ yellow; neotropical. *Trypethelium tolimense* Lücking, Moncada & M. Gut.
- Z6b Only thallus UV+ yellow; pseudostroma UV-. 7
- Z7a Ascospores muriform; Brazil. *Trypethelium muriforme* Aptroot & M.F. Souza
- Z7b Ascospores only transversely septate; Brazil. *Trypethelium endoflavum* Aptroot
- Z8a Pseudostromata lacking pigment, prominent; ascospores (65–)83–97 × 11–15 µm, 13–19-septate; Brazil. *Trypethelium regnellii* Malme
- Z8b Pseudostromata internally with yellow pigment; neotropical. *Trypethelium tolimense* Lücking, B. Mocada & M.C. Gut.
- Z9a Pseudostromata inside and/or outside with yellow or orange pigments (sometimes thin and not readily visible). 10
- Z9b Pseudostromata lacking pigments. 15
- Z10a Ostiolar areas very broad, disc-shaped or fused in lobate fashion; pigments usually sparse. 11
- Z10b Ostiolar areas narrow, visible as small dots; pigments conspicuous 13
- Z11a Ostiolar areas disc-shaped, uniformly brown-black, marginally with small, black papillae arranged in circular fashion. *Trypethelium krempehuberi* Makhija & Patw.
- Z11b Ostiolar areas fused in lobate fashion, contrasting with a bordering whitish rim, without marginal papillae. 12
- Z12a Ostiolar area centrally with dark red pigment in addition to orange pigment bordering the ostiolar lobes; ostioles sometimes eccentric and fused; ascospores 65–100 µm long; neotropical. *Trypethelium astroideum* Flakus & Aptroot

- Z12b Ostiolar area only with bordering orange pigment; ostioles always apical; ascospores 55–75 μm long; pantropical. *Trypethelium platystomum* Mont.
- Z13a Pseudostromata immersed to erumpent; ascospores 37–42 μm long, 7–9-septate; Vietnam. *Trypethelium infraeluteriae* Aptroot & Gueidan
- Z13b Pseudostromata prominent to sessile; ascospores (37–)42–85 μm long, 9–17-septate. 14
- Z14a Pseudostromata inside and/or outside with yellow pigment; ascospores 37–52 \times 8–11 μm , 9–13-septate; pantropical. *Trypethelium eluteriae* Spreng.
- Z14b Pseudostromata inside and/or outside with orange-yellow pigment; ascospores 60–85 \times 11–12 μm , 13–16-septate; pantropical. *Trypethelium subeluteriae* Makhija & Patw.
- Z15a Pseudostromata immersed-erumpent, with ascomata clearly distinguished and individually protruding, brown-black. 16
- Z15b Pseudostromata erumpent to prominent, with ascomata completely immersed in the pseudostromata and not individually protruding, with whitish ostiolar spot or ostiolar area surrounded by irregular whitish to cream-coloured rim or pseudostromata entirely whitish. 17
- Z16a Thallus verrucose-rugose; individual ascomata with flattened top; ascospores 45–55 \times 6–8 μm , 12–17-septate; palaeotropical. *Trypethelium plicatorimosum* Makhija & Patw.
- Z16b Thallus smooth to uneven; individual ascomata with convex top; ascospores 21–40 \times 9–15 μm , 5–11-septate; pantropical. *Trypethelium inaequale* Fée
- Z17a Pseudostromata erumpent to prominent, entirely white or cream-coloured; hamathecium clear. 18
- Z17b Pseudostromata prominent to sessile, grey to black with contrasting, white ostiolar spots; hamathecium interspersed. 19
- Z18a Pseudostromata entirely white, with small, dark ostiolar spots; ascospores 30–60 μm long, 5–9(–13)-septate; palaeotropical. *Trypethelium epileucodes* Nyl.
- Z18b Pseudostromata cream-coloured, with larger, often angular-lobate ostiolar spots; ascospores 58–78 μm long, 9–15-septate; Brazil. *Trypethelium foveolatum* Müll. Arg.
- Z19a Pseudostromata sessile with constricted base, upper part black, sharply contrasting with white ostiolar spots; ascospores 80–110 \times 14–18 μm , 13–19-septate; Africa. *Trypethelium sphaerocephalum* (Vain.) Zahlbr.

Z19b Pseudostromata prominent to sessile, grey to black with contrasting, white ostiolar spots; hamathecium inspersed; neotropical. *Trypethelium ornatum* Müll. Arg.

Ω Key of . *Viridothelium* species

Ω1a Ascospores muriform; thallus and ascomata strongly contrasting in colour, with thallus brownish and ascomata white with black ostiolar areas; neotropical.

Viridothelium tricolor Lücking, M.P. Nelsen & N. Salazar

Ω1b Ascospores transversely septate; thallus and ascomata not strongly contrasting in colour, more or less olive-green to yellowish brown with ascomata sometimes blackish. 2

Ω2a Ascospores 5- or more septate. 3

Ω2b Ascospores 3-septate. 10

Ω3a Ostioles eccentric. 4

Ω3b Ostioles apical. 6

Ω4a Ascospores $75\text{--}98 \times 17\text{--}20 \mu\text{m}$, 15–19-septate; Solomon Islands. *Viridothelium solomonense* Aptroot

Ω4b Ascospores $90\text{--}152 \times 22\text{--}33 \mu\text{m}$. 5

Ω5a Ascospores $99\text{--}152 \times 26\text{--}33 \mu\text{m}$, (9–)15–17-septate; India. *Viridothelium megaspermum* (Makhija & Patw.) Aptroot & Lücking

Ω5b Ascospores $90\text{--}100 \times 22\text{--}26 \mu\text{m}$, 15–21-septate; Fiji. *Viridothelium vonkonratii* Lücking, Naksuwankul & Lumbsch

Ω6a Hamathecium inspersed; ascospores $60\text{--}75 \times 12\text{--}17 \mu\text{m}$, 12–14-septate; Papua New Guinea. *Viridothelium inspersum* Aptroot

Ω6b Hamathecium clear. 7

Ω7a Ascospores 90–150 μm long. 8

Ω7b Ascospores 38–80 μm long. 9

Ω8a Ascospores $90\text{--}105 \times 12\text{--}16 \mu\text{m}$, (9–)13–17-septate; eastern palaeotropical. *Viridothelium indutum* (Stirt.) Aptroot & Lücking

Ω8b Ascospores $100\text{--}150 \times 18\text{--}23 \mu\text{m}$, 17–25-septate; Malaysia. *Viridothelium kinabaluense* Aptroot

Ω9a Ascospores $38\text{--}52 \times 7\text{--}10 \mu\text{m}$; temperate America and Asia. *Viridothelium virens* (Tuck. ex Michener) Lücking, M.P. Nelsen & Aptroot

-
- Ω9b Ascospores $50\text{--}80 \times 8\text{--}15 \mu\text{m}$; temperate Asia. *Viridothelium cinereoglaucescens* (Vain.) Lücking, M.P. Nelsen & Aptroot
- Ω10a Thallus and ascomata thinly furnished with orange pigment; Brazil. *Viridothelium leptoseptatum* Aptroot & M. Cáceres
- Ω10b Thallus and ascomata lacking pigment. 11
- Ω11a Ascomata mostly single, immersed to erumpent; Brazil. *Viridothelium sinuosogelatinosum* Aptroot & M. Cáceres
- Ω11b Ascomata in dense, raised groups; Brazil. *Viridothelium ustulatum* M. Cáceres & Aptroot